

The Poincaré-Hopf Theorem for relative braid classes

S. Munaò, R.C. Vandervorst*

March 2, 2013

Abstract

Braid Floer homology is an invariant of proper relative braid classes [12]. Closed integral curves of 1-periodic Hamiltonian vector fields on the 2-disc may be regarded as braids. If the Braid Floer homology of associated proper relative braid classes is non-trivial, then additional closed integral curves of the Hamiltonian equations are forced via a Morse type theory. In this article we show that certain information contained in the braid Floer homology — the Euler-Floer characteristic — also forces closed integral curves and periodic points of arbitrary vector fields and diffeomorphisms and leads to a Poincaré-Hopf type Theorem. The Euler-Floer characteristic for any proper relative braid class can be computed via a finite cube complex that serves as a model for the given braid class. The results in this paper are restricted to the 2-disc, but can be extended to two-dimensional surfaces (with or without boundary).

1 Introduction

Let $\mathbb{D}^2 \subset \mathbb{R}^2$ denote the standard (closed) 2-disc in the plane with coordinates $x = (p, q)$, i.e. $\mathbb{D}^2 := \{(p, q) \in \mathbb{R}^2 : p^2 + q^2 \leq 1\}$, and let $X(x, t)$ be a smooth 1-periodic vector field on $\mathbb{D}^2$, i.e. $X(x, t+1) = X(x, t)$ for all $x \in \mathbb{D}^2$ and $t \in \mathbb{R}$. The vector field X is tangent to the boundary $\partial\mathbb{D}^2$, i.e. $X(x, t) \cdot \nu = 0$ for all $x \in \partial\mathbb{D}^2$, where ν the outward unit normal on $\partial\mathbb{D}^2$. The set of vector fields satisfying these hypotheses is denoted by $\mathcal{F}_{\parallel}(\mathbb{D}^2 \times \mathbb{R}/\mathbb{Z})$. *Closed integral curves* $x(t)$ of X are integral curves¹ of X for which $x(t + \ell) = x(t)$ for some $\ell \in \mathbb{N}$. Every integral curve of X with minimal period ℓ defines a closed loop in the configuration space $\mathbf{C}_{\ell}(\mathbb{D}^2)$ of ℓ unordered distinct points. A collection of distinct closed integral curves with periods ℓ_j defines

*Department of Mathematics, VU University Amsterdam, The Netherlands.

¹Integral curves of X are smooth functions $x : \mathbb{R} \rightarrow \mathbb{D}^2 \subset \mathbb{R}^2$ that satisfy the differential equation $x' = X(x, t)$.

a closed loop in $\mathbf{C}_m(\mathbb{D}^2)$, with $m = \sum_j \ell_j$. As curves in the cylinder $\mathbb{D}^2 \times [0, 1]$ such a collection of integral curves represents a geometric braid which corresponds to a unique word $\beta_y \in \mathbf{B}_m$, modulo conjugacy and full twists:

$$\beta_y \sim \beta_y \Delta^{2k} \sim \Delta^{2k} \beta_y, \quad (1.1)$$

where Δ^2 is a full positive twist and $\mathbf{B}_m$ is the Artin Braid group on m strands.

Let y be a geometric braid consisting of closed integral curves of X , which will be referred to as a *skeleton*. The curves $y^i(t)$, $i = 1, \dots, m$ satisfy the periodicity condition $y(0) = y(1)$ as point sets, i.e. $y^i(0) = y^{\sigma(i)}(1)$ for some permutation $\sigma \in S_m$. In the configuration space $\mathbf{C}_{n+m}(\mathbb{D}^2)^2$ we consider closed loops of the form $x \text{ rel } y := \{x^1(t), \dots, x^n(t), y^1(t), \dots, y^m(t)\}$. The path component of $x \text{ rel } y$ of closed loops in $\mathcal{L}\mathbf{C}_{n+m}(\mathbb{D}^2)$ is denoted by $[x \text{ rel } y]$ and is called a *relative braid class*. The loops $x' \text{ rel } y' \in [x \text{ rel } y]$, keeping y' fixed, is denoted by $[x'] \text{ rel } y'$ and is called a *fiber*. Relative braid classes are path components of braids which have at least two components and the components are labeled into two groups: x and y . The intertwining of x and y defines various different braid classes. A relative braid class $[x \text{ rel } y]$ in $\mathbb{D}^2$ is *proper* if components $x_c \subset x$ cannot be deformed onto (i) the boundary $\partial\mathbb{D}^2$, (ii) itself,³ or other components $x'_c \subset x$, or (iii) components in $y_c \subset y$, see [12] for details. In this paper we are mainly concerned with relative braids for which x has only one strand. To proper relative braid classes $[x \text{ rel } y]$ one can assign the invariants $\text{HB}_*([x \text{ rel } y])$, with coefficients in $\mathbb{Z}_2$, called *Braid Floer homology*. In the following subsection we will briefly explain the construction of the invariants $\text{HB}_*([x \text{ rel } y])$ in case that x consists of one single strand. See [12] for more details on Braid Floer homology.

1.1 A brief summary of Braid Floer homology

Fix a Hamiltonian vector field X_H in $\mathcal{F}_{||}(\mathbb{D}^2 \times \mathbb{R}/\mathbb{Z})$ of the form $X_H(x, t) = J\nabla H(x, t)$, where

$$J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

and H is a Hamiltonian function with the properties:

- (i) $H \in C^\infty(\mathbb{D}^2 \times \mathbb{R}/\mathbb{Z}; \mathbb{R})$;

²The space of continuous mapping $\mathbb{R}/\mathbb{Z} \rightarrow X$, with X a topological space, is called the free loop space of X and is denoted by $\mathcal{L}X$.

³This condition is separated into two cases: (i) a component in x cannot be not deformed into a single strand, or (ii) if a component in x can be deformed into a single strand, then the latter necessarily intersects y or a different component in x .

(ii) $H(x, t)|_{x \in \partial \mathbb{D}^2} = 0$, for all $t \in \mathbb{R}/\mathbb{Z}$.

For closed integral curves of X_H of period 1 we define the Hamilton action

$$\mathcal{A}_H(x) = \int_0^1 \frac{1}{2} Jx \cdot x_t - H(x, t) dt,$$

Critical points of the action functional $\mathcal{A}_H$ are in one-to-one correspondence with closed integral curves of period 1. Assume that $y = \{y^j(t)\}$ is a collection of closed integral curves of the Hamilton vector field X_H , i.e. periodic solutions of the $y_t^j = X_H(y^j, t)$. Consider a proper relative braid class $[x] \text{ rel } y$, with x 1-periodic and seek closed integral curves $x \text{ rel } y$ in $[x] \text{ rel } y$. The set of critical points of $\mathcal{A}_H$ in $[x] \text{ rel } y$ is denoted by $\text{Crit}_{\mathcal{A}_H}([x] \text{ rel } y)$. In order to understand the set $\text{Crit}_{\mathcal{A}_H}([x] \text{ rel } y)$ we consider the negative L^2 -gradient flow of $\mathcal{A}_H$. The L^2 -gradient flow $u_s = -\nabla_{L^2} \mathcal{A}_H(u)$ yields the Cauchy-Riemann equations

$$u_s(s, t) + Ju_t(s, t) + \nabla H(u(s, t), t) = 0,$$

for which the stationary solutions $u(s, t) = x(t)$ are the critical points of $\mathcal{A}_H$.

To a braid y one can assign an integer $\text{Cross}(y)$ which counts the number of crossings (with sign) of strands in the standard planar projection. In the case of a relative braid $x \text{ rel } y$ the number $\text{Cross}(x \text{ rel } y)$ is an invariant of the relative braid class $[x \text{ rel } y]$. In [12] a monotonicity lemma is proved, which states that along solutions $u(s, t)$ of the nonlinear Cauchy-Riemann equations, the number $\text{Cross}(u(s, \cdot) \text{ rel } y)$ is non-increasing (the jumps correspond to ‘singular braids’, i.e. ‘braids’ for which intersections occur). As a consequence an isolation property for proper relative braid classes exists: the set bounded solutions of the Cauchy-Riemann equations in a proper braid class fiber $[x] \text{ rel } y$, denoted by $\mathcal{M}([x] \text{ rel } y; H)$, is compact and isolated with respect to the topology of uniform convergence on compact subsets of $\mathbb{R}^2$. These facts provide all the ingredients to follow Floer’s approach towards Morse Theory for the Hamiltonian action [7]. For generic Hamiltonians which satisfy (i) and (ii) above and for which y is a skeleton, the critical points in $[x] \text{ rel } y$ of the action $\mathcal{A}_H$ are non-degenerate and the set of connecting orbits $\mathcal{M}_{x_-, x_+}([x] \text{ rel } y; H)$ are smooth finite dimensional manifolds. To critical in $\text{Crit}_{\mathcal{A}_H}([x] \text{ rel } y)$ we assign a relative index $\mu^{CZ}(x)$ (the Conley-Zehnder index) and

$$\dim \mathcal{M}_{x_-, x_+}([x] \text{ rel } y; H) = \mu^{CZ}(x_-) - \mu^{CZ}(x_+).$$

Define the free abelian groups C_k over the critical points of index k , with coefficients in $\mathbb{Z}_2$, i.e.

$$C_k([x] \text{ rel } y; H) := \bigoplus_{\substack{x \in \text{Crit}_{\mathcal{A}_H}([x] \text{ rel } y), \\ \mu(x) = k}} \mathbb{Z}_2 \langle x \rangle,$$

and the boundary operator

$$\partial_k = \partial_k([x] \text{ rel } y; H) : C_k \rightarrow C_{k-1},$$

which counts the number of orbits (modulo 2) between critical points of index k and $k - 1$ respectively. Analysis of the spaces $\mathcal{M}_{x_-, x_+}([x] \text{ rel } y; H)$ reveals that (C_*, ∂_*) is a chain complex, and its (Floer) homology is denoted by $\text{HB}_*([x] \text{ rel } y; H)$. Different choices of H yields isomorphic Floer homologies and

$$\text{HB}_*([x] \text{ rel } y) = \varprojlim \text{HB}_*([x] \text{ rel } y; H),$$

where the inverse limit is defined with respect to the canonical isomorphisms $a_k(H, H') : \text{HB}_k([x] \text{ rel } y, H) \rightarrow \text{HB}_k([x] \text{ rel } y, H')$. Some properties are:

- (i) the groups $\text{HB}_k([x] \text{ rel } y)$ are defined for all $k \in \mathbb{Z}$ and are finite, i.e. $\mathbb{Z}_2^d$ for some $d \geq 0$;
- (ii) the groups $\text{HB}_k([x] \text{ rel } y)$ are invariants for the fibers in the same relative braid class $[x \text{ rel } y]$, i.e. if $x \text{ rel } y \sim x' \text{ rel } y'$, then $\text{HB}_k([x] \text{ rel } y) \cong \text{HB}_k([x'] \text{ rel } y')$. For this reason we will write $\text{HB}_*([x \text{ rel } y])$;
- (iii) if $(x \text{ rel } y) \cdot \Delta^{2\ell}$ denotes composition with ℓ full twists, then $\text{HB}_k([(x \text{ rel } y) \cdot \Delta^{2\ell}]) \cong \text{HB}_{k-2\ell}([x \text{ rel } y])$.

1.2 The Euler-Floer characteristic and the Poincaré-Hopf Formula

Braid Floer homology is an invariant of conjugacy classes in $\mathbf{B}_{n+m}$ and can be computed from purely topological data. The *Euler-Floer characteristic* of $\text{HB}_*([x \text{ rel } y])$ is defined as follows:

$$\chi(x \text{ rel } y) = \sum_{k \in \mathbb{Z}} (-1)^k \dim \text{HB}_k([x \text{ rel } y]). \quad (1.2)$$

In Section 7 we show that the Euler-Floer characteristic of $\text{HB}_*([x \text{ rel } y])$ can be computed from a finite cube complex which serves as a model for the braid class.

A 1-periodic function $x \in C^1(\mathbb{R}/\mathbb{Z})$ is an *isolated* closed integral curve of X if there exists an $\epsilon > 0$ such that x is the only solution of the differential equation

$$\mathcal{E}(x(t)) = \frac{dx}{dt}(t) - X(x(t), t), \quad (1.3)$$

in $B_\epsilon(x) \subset C^1(\mathbb{R}/\mathbb{Z})$. For isolated, and in particular non-degenerate closed integral curves we can define an *index* as follows. Let $\Theta \in \text{M}_{2 \times 2}(\mathbb{R})$ be any matrix satisfying $\sigma(\Theta) \cap 2\pi k i \mathbb{R} = \emptyset$, for all $k \in \mathbb{Z}$ and let $\eta \mapsto R(t; \eta)$ be a curve in $C^\infty(\mathbb{R}/\mathbb{Z}; \text{M}_{2 \times 2}(\mathbb{R}))$,

with $R(t; 0) = \Theta$ and $R(t; 1) = D_x X(x(t), t)$ — the linearization of X at $x(t)$. Then $\eta \mapsto F(\eta) = \frac{d}{dt} - R(t; \eta)$ defines a curve in $\text{Fred}_0(C^1, C^0)$, where we denote by $\text{Fred}_0(C^1, C^0)$ the space of Fredholm operators of index 0 between C^1 and C^0 . Denote by $\Sigma \subset \text{Fred}_0(C^1, C^0)$ the set of non-invertible operators and by $\Sigma_1 \subset \Sigma$ the non-invertible operators with a 1-dimensional kernel. If the end points of F are invertible one can choose the path $\eta \mapsto R(t; \eta)$ such that $F(\eta)$ intersects Σ in Σ_1 and all intersections are transverse. If $\gamma = \#$ intersections of $F(\eta)$ with Σ_1 , then

$$\iota(x) = -\text{sgn}(\det(\Theta))(-1)^\gamma. \quad (1.4)$$

This definition is independent of the choice of Θ , see Section 6.

The above definition can be expressed in terms of the Leray-Schauder degree. Let $M \in \text{GL}(C^0, C^1)$ be any isomorphism such that $\Phi_M(x) := M\mathcal{E}(x)$ is of the form ‘identity + compact’. Then the index of an isolated closed integral curve is given by

$$\iota(x) = -\text{sgn}(\det(\Theta))(-1)^{\beta_M(\Theta)} \deg_{LS}(\Phi_M, B_\epsilon(x), 0). \quad (1.5)$$

where $\beta_M(\Theta)$ is the number of negative eigenvalues of $M \frac{d}{dt} - M\Theta$ counted with multiplicity. The latter definition holds for both non-degenerate and isolated 1-periodic closed integral curves of X . In Section 6 we show that the two expressions for the index are the same and we show that they are independent of the choices of M and Θ .

Theorem 1.1 (Poincaré-Hopf Formula). *Let y be a skeleton of closed integral curves of a vector field $X \in \mathcal{F}_\parallel(\mathbb{D}^2 \times \mathbb{R}/\mathbb{Z})$ and let $[x \text{ rel } y]$ be a proper relative braid class. Suppose that all 1-periodic closed integral curves of X are isolated, then for all closed integral curves $x_0 \text{ rel } y$ in $[x_0] \text{ rel } y$ it holds that*

$$\sum_{x_0} \iota(x_0) = \chi(x \text{ rel } y). \quad (1.6)$$

The index formula can be used to obtain existence results for closed integral curves in proper relative braid classes.

Theorem 1.2. *Let y be a skeleton of closed integral curves of a vector field $X \in \mathcal{F}_\parallel(\mathbb{D}^2 \times \mathbb{R}/\mathbb{Z})$ and let $[x \text{ rel } y]$ be a proper relative braid class. If $\chi(x \text{ rel } y) \neq 0$, then there exist closed integral curves $x_0 \text{ rel } y$ in $[x] \text{ rel } y$.*

The analogue of Theorem 1.1 can also be proved for relative braid class $[x \text{ rel } y]$ in $\mathbf{C}_{n+m}(\mathbb{D}^2)$. Our theory also provides detailed information about the linking of solutions. In Section 8 we give various examples and compute the Euler-Floer characteristic. This does not provide a procedure for computing the braid Floer homology.

Remark 1.1. In this paper Theorem 1.1 is proved using the standard Leray-Schauder degree theory in combination with the theory of spectral flow and parity for operators on Hilbert spaces. The Leray-Schauder degree is related to the Euler characteristic of Braid Floer homology. An other approach is the use the degree theory developed by Fitzpatrick et al. [5].

1.3 Discretization and computability

The second part of the paper deals with the computability of the Euler-Floer characteristic. This is obtained through a finite dimensional model. A model is constructed in three steps:

- (i) compose $x \text{ rel } y$ with $\ell \geq 0$ full twists Δ^2 , such that $(x \text{ rel } y) \cdot \Delta^{2\ell}$ is isotopic to a positive braid $x^+ \text{ rel } y^+$;
- (ii) relative braids $x^+ \text{ rel } y^+$ are isotopic to Legendrian braids $x_L \text{ rel } y_L$ on $\mathbb{R}^2$, i.e. braids which have the form $x_L = (q_t, q)$ and $y_L = (Q_t, Q)$, where $q = \pi_2 x$ and $Q = \pi_2 y$, and π_2 the projection onto the q -coordinate;
- (iii) discretize q and $Q = \{Q^j\}$ to $q_d = \{q_i\}$, with $q_i = q(i/d)$, $i = 0, \dots, d$ and $Q_D = \{Q_D^j\}$, with $Q_D^j = \{Q_i^j\}$ and $Q_i^j = Q^j(i/d)$ respectively, and consider the piecewise linear interpolations connecting the *anchor* points q_i and Q_i^j for $i = 0, \dots, d$. A discretization $q_D \text{ rel } Q_D$ is *admissible* if the linear interpolation is isotopic to $q \text{ rel } Q$. All such discretization form the discrete relative braid class $[q_D \text{ rel } Q_D]$, for which each fiber is a finite cube complex.

Remark 1.2. If the number of discretization points is not large enough, then the discretization may not be admissible and therefore not capture the topology of the braid. See [8] and Section 7.4 for more details.

For $d > 0$ large enough there exists an admissible discretization $q_D \text{ rel } Q_D$ for any Legendrian representative $x_L \text{ rel } y_L \in [x \text{ rel } y]$ and thus an associated discrete relative braid class $[q_D \text{ rel } Q_D]$. In [8] an invariant for discrete braid classes was introduced. Let $[q_D] \text{ rel } Q_D$ denote a fiber in $[q_D \text{ rel } Q_D]$, which is a cube complex with a finite number of connected components and their closures are denoted by N_j . The faces of the hypercubes N_j can be co-oriented in direction of decreasing the number of crossing in $q_D \text{ rel } Q_D$, and we define N_j^- as the closure of the set of faces with outward pointing co-orientation. The sets N_j^- are called *exit sets*. The invariant for a fiber is given by

$$\text{HC}_*([q_D] \text{ rel } Q_D) = \bigoplus_j H_*(N_j, N_j^-).$$

This discrete braid invariant is well-defined for any $d > 0$ for which there exist admissible discretizations and is independent of both the particular fiber and the discretization size d . For the associated Euler characteristic we therefore write $\chi(q_D \text{ rel } Q_D)$. The latter is an Euler characteristic of a topological pair. The Euler characteristic of the Braid Floer homology $\chi(x \text{ rel } y)$ can be related to the Euler characteristic of the associated discrete braid class.

Theorem 1.3. *Let $[x \text{ rel } y]$ a proper relative braid class and $\ell \geq 0$ is an integer such that $(x \text{ rel } y) \cdot \Delta^{2\ell}$ is isotopic to a positive braid $x^+ \text{ rel } y^+$. Let $q_D \text{ rel } Q_D$ be an admissible discretization, for some $d > 0$, of a Legendrian representative $x_L \text{ rel } y_L \in [x^+ \text{ rel } y^+]$. Then*

$$\chi(x \text{ rel } y) = \chi(q_D \text{ rel } Q_D^*),$$

where Q_D^* is the augmentation of Q_D by adding the constant strands ± 1 to Q_D .

The idea behind the proof of Theorem 1.3 is to first relate $\chi(x \text{ rel } y)$ to mechanical Lagrangian systems and then use a discretization approach based on the method of broken geodesics. Theorem 1.3 is proved in Section 7. In Section 8 we use the latter to compute the Euler-Floer characteristic for various examples of proper relative braid classes.

1.4 Additional topological properties

In this paper we do not address the question whether the closed integral curves $x \text{ rel } y$ are non-constant, i.e. are not equilibrium points. By considering relative braid classes where x consists of more than one strand one can study non-constant closed integral curves. Braid Floer homology for relative braids with x consisting of n strands is defined in [12]. The ideas in this paper extend to relative braid classes with multi-strand braids x . In Section 8 we give an example of a multi-strand x in $x \text{ rel } y$ and explain how this yields the existence of non-trivial closed integral curves.

The invariant $\chi(q_D \text{ rel } Q_D)$ is a true Euler characteristic and

$$\chi(q_D \text{ rel } Q_D) = \chi([q_D] \text{ rel } Q_D, [q_D]^- \text{ rel } Q_D),$$

where $[q_D]^- \text{ rel } Q_D$ is the exit. A similar characterization does not a priori exist for $[x] \text{ rel } y$. This problem is circumvented by considering Hamiltonian systems and carrying out Floer's approach towards Morse theory (see [7]), by using the isolation property of $[x] \text{ rel } y$. The fact that the Euler characteristic of Floer homology is related to the Euler characteristic of a topological pair indicates that Floer homology is a good substitute for a suitable (co)-homology theory. For more details see Section 7 and Remark 7.1.

Braid Floer homology developed for the 2-disc $\mathbb{D}^2$ can be extended to more general 2-dimensional manifolds. This generalization of Braid Floer homology for 2-dimensional manifolds can then be used to extend the results in this paper to more general surfaces.

Acknowledgment. The authors wish to thank J.B. van den Berg for the many stimulating discussions on the subject of Braid Floer homology.

2 Closed integral curves

Let $X \in \mathcal{F}_\parallel(\mathbb{D}^2 \times \mathbb{R}/\mathbb{Z})$, then closed integral curves of X of period 1 satisfy the differential equation

$$\begin{cases} \frac{dx}{dt} = X(x, t), & x \in \mathbb{D}^2, t \in \mathbb{R}/\mathbb{Z}, \\ x(0) = x(1). \end{cases} \quad (2.1)$$

Consider the unbounded operator $L_\mu : C^1(\mathbb{R}/\mathbb{Z}) \subset C^0(\mathbb{R}/\mathbb{Z}) \rightarrow C^0(\mathbb{R}/\mathbb{Z})$, defined by

$$L_\mu := -J \frac{d}{dt} + \mu, \quad \mu \in \mathbb{R}.$$

The operator is invertible for $\mu \neq 2\pi k, k \in \mathbb{Z}$ and the inverse $L_\mu^{-1} : C^0(\mathbb{R}/\mathbb{Z}) \rightarrow C^0(\mathbb{R}/\mathbb{Z})$ is compact. Transforming Equation (2.1), using L_μ^{-1} , yields the equation $\Phi_\mu(x) = 0$, where

$$\Phi_\mu(x) := x - L_\mu^{-1}(-JX(x, t) + \mu x).$$

If we set

$$K_\mu(x) := L_\mu^{-1}(-JX(x, t) + \mu x),$$

then Φ_μ is of the form $\Phi_\mu(x) = x - K_\mu(x)$, where K_μ is a (non-linear) compact operator on $C^0(\mathbb{R}/\mathbb{Z})$. Since X is a smooth vector field the mapping Φ_μ is a smooth mapping on $C^0(\mathbb{R}/\mathbb{Z})$.

Proposition 2.1. *A function $x \in C^0(\mathbb{R}/\mathbb{Z})$, with $|x(t)| \leq 1$ for all t , is a solution of $\Phi_\mu(x) = 0$ if and only if $x \in C^1(\mathbb{R}/\mathbb{Z})$ and x satisfies Equation (2.1).*

Proof. If $x \in C^1(\mathbb{R}/\mathbb{Z}; \mathbb{D}^2)$ is a solution of Equation (2.1), then $\Phi_\mu(x) = 0$ is obviously satisfied. On the other hand, if $x \in C^0(\mathbb{R}/\mathbb{Z}; \mathbb{D}^2)$ is a zero of Φ_μ , then $x = K_\mu(x) \in C^1(\mathbb{R}/\mathbb{Z})$, since $R(L_\mu^{-1}) \subset C^1(\mathbb{R}/\mathbb{Z})$. Applying L_μ to both sides shows that x satisfies Equation (2.1). $\square$

Note that the zero set $\Phi_\mu^{-1}(0)$ does not depend on the parameter μ . In order to apply the Leray-Schauder degree theory we consider appropriate bounded, open subsets $\Omega \subset C^0(\mathbb{R}/\mathbb{Z})$, which have the property that $\Phi_\mu^{-1}(0) \cap \partial\Omega = \emptyset$. Let $\Omega = [x] \text{ rel } y$, where $[x] \text{ rel } y$ is a proper relative braid fiber, and $y = \{y^1, \dots, y^m\}$ is a skeleton of closed integral curves for the vector field X .

Proposition 2.2. *Let $[x \text{ rel } y]$ be a proper relative braid class and let $\Omega = [x] \text{ rel } y$ be the fiber given by y . Then, there exists an $0 < r < 1$ such that*

$$|x(t)| < r, \text{ and } |x(t) - y^j(t)| > 1 - r, \quad \forall j = 1, \dots, m, \quad \forall t \in \mathbb{R},$$

and for all $x \in \Phi_\mu^{-1}(0) \cap \Omega = \{x \in \Omega \mid x = K_\mu(x)\}$.

Proof. Since $\Omega \subset C^0(\mathbb{R}/\mathbb{Z})$ is a bounded set and K_μ is compact, the solution set $\Phi_\mu^{-1}(0) \cap \Omega$ is compact. Indeed, let $x_n = K_\mu(x_n)$ be a sequence in $\Phi_\mu^{-1}(0) \cap \Omega$, then $K_\mu(x_{n_k}) \rightarrow x$, and thus $x_{n_k} \rightarrow x$, which, by continuity, implies that $K_\mu(x_{n_k}) \rightarrow K_\mu(x)$, and thus $x \in \Phi_\mu^{-1}(0) \cap \Omega$.

Let $x_n \in \Phi_\mu^{-1}(0) \cap \Omega$ and assume that such an $0 < r < 1$ does not exist. Then, by the compactness of $\Phi_\mu^{-1}(0) \cap \Omega$, there is a subsequence $x_{n_k} \rightarrow x$ such that one, or both of the following two possibilities hold: (i) $|x(t_0)| = 1$ for some t_0 . By the uniqueness of solutions of Equation (2.1) and the invariance of the boundary $\partial\mathbb{D}^2$ ($X(x, t)$ is tangent to the boundary), $|x(t)| = 1$ for all t , which is impossible since $[x] \text{ rel } y$ is proper; (ii) $x(t_0) = y^j(t_0)$ for some t_0 and some j . As before, by the uniqueness of solutions of Equation (2.1), then $x(t) = y^j(t)$ for all t , which again contradicts the fact that $[x] \text{ rel } y$ is proper. $\square$

By Proposition 2.2 the Leray-Schauder degree $\deg_{LS}(\Phi_\mu, \Omega, 0)$ is well-defined. Consider the Hamiltonian vector field

$$X_H = J\nabla H, \quad J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad (2.2)$$

where $H(x, t)$ is a smooth Hamiltonian such that $X_H \in \mathcal{F}_\parallel(\mathbb{D}^2 \times \mathbb{R}/\mathbb{Z})$ and y is a skeleton for X_H . Such a Hamiltonian can always be constructed, see [12], and the class of such Hamiltonians will be denote by $\mathcal{H}_\parallel(y)$. Since y is a skeleton for both X and X_H , it is a skeleton for the linear homotopy $X_\alpha = (1 - \alpha)X + \alpha X_H$, $\alpha \in [0, 1]$. Associated with the homotopy X_α of vector fields we define the homotopy

$$\Phi_{\mu, \alpha}(x) := x - L_\mu^{-1}(-JX_\alpha(x, t) + \mu x) = x - K_{\mu, \alpha}(x), \quad \alpha \in [0, 1],$$

with $K_{\mu, \alpha}(x) = L_\mu^{-1}(-JX_\alpha(x, t) + \mu x)$. Proposition 2.2 applies for all $\alpha \in [0, 1]$, i.e. by compactness there exists a uniform $0 < r < 1$ such that

$$|x(t)| < r, \text{ and } |x(t) - y^j(t)| > 1 - r,$$

for all $t \in \mathbb{R}$, for all j and for all $x \in \Phi_{\mu,\alpha}^{-1}(0) \cap \Omega = \{x \in \Omega \mid x = K_{\mu,\alpha}(x)\}$ and all $\alpha \in [0, 1]$. By the homotopy invariance of the Leray-Schauder degree we have

$$\deg_{LS}(\Phi_\mu, \Omega, 0) = \deg_{LS}(\Phi_{\mu,\alpha}, \Omega, 0) = \deg_{LS}(\Phi_{\mu,H}, \Omega, 0), \quad (2.3)$$

where $\Phi_{\mu,0} = \Phi_\mu$ and $\Phi_{\mu,1} = \Phi_{\mu,H}$. Note that the zeroes of $\Phi_{\mu,H}$ correspond to critical point of the functional

$$\mathcal{A}_H(x) = \int_0^1 \frac{1}{2} Jx \cdot x_t - H(x, t) dt, \quad (2.4)$$

and are denoted by $\text{Crit}_{\mathcal{A}_H}([x] \text{ rel } y)$. In [12] invariants are defined which provide information about $\Phi_{\mu,H}^{-1}(0) \cap \Omega = \text{Crit}_{\mathcal{A}_H}([x] \text{ rel } y)$ and thus $\deg_{LS}(\Phi_{\mu,H}, \Omega, 0)$. These invariants are the Braid Floer homology groups $\text{HB}_*([x] \text{ rel } y)$ as explained in the introduction. In the next section we examine spectral properties of the solutions of $\Phi_{\mu,\alpha}^{-1}(0) \cap \Omega$ in order to compute $\deg_{LS}(\Phi_{\mu,H}, \Omega, 0)$ and thus $\deg_{LS}(\Phi_\mu, \Omega, 0)$.

Remark 2.1. There is obviously more room for choosing appropriate operators L_μ and therefore functions Φ_μ . In Section 6 this issue will be discussed in more detail.

3 Parity, Spectral flow and the Leray-Schauder degree

The Leray-Schauder degree of an isolated zero x of $\Phi_\mu(x) = 0$ is called the local degree. A zero $x \in \Phi_\mu^{-1}(0)$ is non-degenerate if $1 \notin \sigma(D_x K_\mu(x))$, where $D_x K_\mu(x) : C^0(\mathbb{R}/\mathbb{Z}) \rightarrow C^0(\mathbb{R}/\mathbb{Z})$ is the (compact) linearization at x and is given by $D_x K_\mu(x) = L_\mu^{-1}(-JD_x X(x, t) + \mu)$. If x is a non-degenerate zero, then it is an isolated zero and the degree can be determined from spectral information.

Proposition 3.1. *Let $x \in C^0(\mathbb{R}/\mathbb{Z})$ be a non-degenerate zero of Φ_μ and let $\epsilon > 0$ be sufficiently small such that $B_\epsilon(x) = \{\tilde{x} \in C^0(\mathbb{R}/\mathbb{Z}) \mid |\tilde{x}(t) - x(t)| < \epsilon, \forall t\}$ is a neighborhood in which x is the only zero. Then*

$$\deg_{LS}(\Phi_\mu, B_\epsilon(x), 0) = \deg_{LS}(\text{Id} - D_x K_\mu(x), B_\epsilon(x), 0) = (-1)^{\beta_\mu(x)}$$

where

$$\beta_\mu(x) = \sum_{\sigma_j > 1, \sigma_j \in \sigma(D_x K_\mu(x))} \beta_j, \quad \beta_j = \dim \left(\bigcup_{i=1}^{\infty} \ker(\sigma_j \text{Id} - D_x K_\mu(x))^i \right),$$

which will be referred to as the Morse index of x , or alternatively the Morse index of linearized operator $D_x \Phi_\mu(x)$.

Proof. See [9]. □

The functions $\Phi_{\mu,\alpha}(x) = x - K_{\mu,\alpha}(x)$ are of the form ‘identity + compact’ and Proposition 3.1 can be applied to non-degenerate zeroes of $\Phi_{\mu,\alpha}(x) = 0$. If we choose the Hamiltonian $H \in \mathcal{H}_{\parallel}^{\text{reg}}(y)$ ‘generically’, then the zeroes of $\Phi_{\mu,H}$ are non-degenerate, i.e. $1 \notin \sigma(D_x K_{\mu,H}(x))$, where $D_x K_{\mu,H}(x) = D_x K_{\mu,1}(x)$. By compactness there are only finitely many zeroes in a fiber $\Omega = [x] \text{ rel } y$.

Lemma 3.1. *Let $x \in \Phi_{\mu,H}^{-1}(0) \cap \Omega$. Then following criteria for non-degeneracy are equivalent:*

- (i) $1 \notin \sigma(D_x K_{\mu,H}(x))$;
- (ii) the operator $B = -J \frac{d}{dt} - D_x^2 H(x(t), t)$ is invertible;
- (iii) let $\Psi(t)$ be defined by $B\Psi(t) = 0$, $\Psi(0) = \text{Id}$, then $\det(\Psi(1) - \text{Id}) \neq 0$.

Proof. A function ψ satisfies $D_x K_{\mu,H}(x)\psi = \psi$ if and only if $B\psi = 0$, which shows the equivalence between (i) and (ii). The equivalence between (ii) and (iii) is proved in [12]. $\square$

The generic choice of H follows from Proposition 7.1 in [12] based on criterion (iii). Hamiltonians for which the zeroes of $\Phi_{\mu,H}$ are non-degenerate are denoted by $\mathcal{H}_{\parallel}^{\text{reg}}(y)$. Note that *no* genericity is needed for $\alpha \in [0, 1]$! For the Leray-Schauder degree this yields

$$\deg_{LS}(\Phi_{\mu,\alpha}, \Omega, 0) = \deg_{LS}(\Phi_{\mu,H}, \Omega, 0) = \sum_{x \in \text{Crit}_{\mathcal{H}}([x] \text{ rel } y)} (-1)^{\beta_{\mu,H}(x)}, \quad (3.1)$$

for all $\alpha \in [0, 1]$ and where $\beta_{\mu,H}(x)$ is the Morse index of $\text{Id} - D_x K_{\mu,H}(x)$.

The goal is to determine the Leray-Schauder degree $\deg_{LS}(\Phi_{\mu}, \Omega, 0)$ from information contained in the Braid Floer homology groups $\text{HB}_*([x] \text{ rel } y)$. In order to do so we examine the Hamiltonian case. In the Hamiltonian case the linearized operator $D_x \Phi_{\mu,H}(x)$ is given by

$$A := D_x \Phi_{\mu,H}(x) = \text{Id} - D_x K_{\mu,H}(x) = \text{Id} - L_{\mu}^{-1}(D_x^2 H(x(t), t) + \mu),$$

which is a bounded operator on $C^0(\mathbb{R}/\mathbb{Z})$. The operator A extends to a bounded operator on $L^2(\mathbb{R}/\mathbb{Z})$. Consider the path $\eta \mapsto A(\eta)$, $\eta \in I = [0, 1]$, given by

$$A(\eta) = \text{Id} - L_{\mu}^{-1}(S(t; \eta) + \mu) = \text{Id} - T_{\mu}(\eta), \quad (3.2)$$

where $S(t; \eta)$ a smooth family of symmetric matrices and $T_{\mu}(\eta) = L_{\mu}^{-1}(S(t; \eta) + \mu)$. The endpoints satisfy

$$S(t; 0) = \theta \text{Id}, \quad S(t; 1) = D_x^2 H(x(t), t),$$

with $\theta \neq 2\pi k$, for some $k \in \mathbb{Z}$ and $D_x^2 H(x(t), t)$ is the Hessian of H at a critical point in $\text{Crit}_{\mathcal{A}_H}([x] \text{ rel } y)$. The path of $\eta \mapsto A(\eta)$ is a path bounded linear Fredholm operators on $L^2(\mathbb{R}/\mathbb{Z})$ of Fredholm index 0, which are compact perturbations of the identity and whose endpoints are invertible.

Lemma 3.2. *The path $\eta \mapsto A(\eta)$ defined in (3.2) is a smooth path of bounded linear Fredholm operators in $H^s(\mathbb{R}/\mathbb{Z})$ of index 0, with invertible endpoints.*

Proof. By the smoothness of $S(t; \eta)$ we have that $\|S(t; \eta)x\|_{H^m} \leq C\|x\|_{H^m}$, for any $x \in H^m(\mathbb{R}/\mathbb{Z})$ and any $m \in \mathbb{N} \cup \{0\}$. By interpolation the same holds for all $x \in H^s(\mathbb{R}/\mathbb{Z})$ and the claim follows from the fact that $L_\mu^{-1} : H^s(\mathbb{R}/\mathbb{Z}) \rightarrow H^{s+1}(\mathbb{R}/\mathbb{Z}) \hookrightarrow H^s(\mathbb{R}/\mathbb{Z})$ is compact. $\square$

3.1 Parity of paths of linear Fredholm operators

Let $\eta \mapsto \Lambda(\eta)$ be a smooth path of bounded linear Fredholm operators of index 0 on a Hilbert space $\mathcal{H}$. A crossing $\eta_0 \in I$ is a number for which the operator $\Lambda(\eta_0)$ is not invertible. A crossing is simple if $\dim \ker \Lambda(\eta_0) = 1$. A path $\eta \mapsto \Lambda(\eta)$ between invertible ends can always be perturbed to have only simple crossings. Such paths are called generic. Following [3–6], we define the *parity* of a generic path $\eta \mapsto \Lambda(\eta)$ by

$$\text{parity}(\Lambda(\eta), I) := \prod_{\ker \Lambda(\eta_0) \neq 0} (-1) = (-1)^{\text{cross}(\Lambda(\eta), I)}, \quad (3.3)$$

where $\text{cross}(\Lambda(\eta), I) = \#\{\eta_0 \in I : \ker \Lambda(\eta_0) \neq 0\}$. The parity is a homotopy invariant with values in $\mathbb{Z}_2$. In [3–6] an alternative characterization of parity is given via the Leray-Schauder degree. For any Fredholm path $\eta \mapsto \Lambda(\eta)$ there exists a path $\eta \mapsto M(\eta)$, called a *parametrix*, such that $\eta \mapsto M(\eta)\Lambda(\eta)$ is of the form ‘identity + compact’. For parity this gives:

$$\text{parity}(\Lambda(\eta), I) = \deg_{LS}(M(0)\Lambda(0)) \cdot \deg_{LS}(M(1)\Lambda(1)),$$

where $\deg_{LS}(M(\eta)\Lambda(\eta)) = \deg_{LS}(M(\eta)\Lambda(\eta), \mathcal{H}, 0)$, for $\eta = 0, 1$, and the expression is independent of the choice of parametrix. The latter extends the above definition to arbitrary paths with invertible endpoints. For a list of properties of parity see [3–6].

Proposition 3.2. *Let $\eta \mapsto A(\eta)$ be the path of bounded linear Fredholm operators on $H^s(\mathbb{R}/\mathbb{Z})$ defined by (3.2). Then*

$$\text{parity}(A(\eta), I) = (-1)^{\beta_{A(0)}} \cdot (-1)^{\beta_{A(1)}} = (-1)^{\beta_{A(0)} - \beta_{A(1)}}. \quad (3.4)$$

where $\beta_{A(0)}$ and $\beta_{A(1)}$ are the Morse indices of $A(0)$ and $A(1)$ respectively.

Proof. For $\eta \mapsto A(\eta)$ the parametrix is the constant path $\eta \mapsto M(\eta) = \text{Id}$. From Proposition 3.1 we derive that

$$\deg_{LS}(A(0)) = (-1)^{\beta_{A(0)}}, \quad \text{and} \quad \deg_{LS}(A(1)) = (-1)^{\beta_{A(1)}},$$

which proves the first part of the formula. Since $\beta(A(0)) - \beta(A(1)) = [\beta(A(0)) + \beta(A(1))] \bmod 2$, the second identity follows. $\square$

Lemma 3.3. *For $\theta > 0$, the Morse index for $A(0)$ is given by $\beta_{A(0)} = 2 \left\lceil \frac{\mu + \theta}{2\pi} \right\rceil$.*

Proof. The eigenvalues of the operator $A(0)$ are given by $\lambda = \frac{-\theta + 2k\pi}{\mu + 2k\pi}$ and all have multiplicity 2. Therefore number of integers k for which $\lambda < 0$ is equal to $\left\lceil \frac{\mu + \theta}{2\pi} \right\rceil$ and consequently $\beta_{A(0)} = 2 \left\lceil \frac{\mu + \theta}{2\pi} \right\rceil$. $\square$

If $x \in \Phi_{\mu,H}^{-1}(0)$ is a non-degenerate zero, then its local degree can be expressed in terms of the parity of $A(\eta)$.

Proposition 3.3. *Let $x \in \Phi_{\mu,H}^{-1}(0)$ be a non-degenerate zero, then*

$$\deg_{LS}(\Phi_{\mu,H}, B_\epsilon(x), 0) = \text{parity}(A(\eta), I), \quad (3.5)$$

where $\eta \mapsto A(\eta)$ is given by (3.2).

Proof. From Proposition 3.1 we have that $\deg_{LS}(\Phi_{\mu,H}, B_\epsilon(x), 0) = (-1)^{\beta_{A(1)}}$ and by Equation (3.4), $\text{parity}(A(\eta), I) = (-1)^{\beta_{A(0)}} \cdot (-1)^{\beta_{A(1)}} = (-1)^{\beta_{A(1)}}$, which completes the proof. $\square$

3.2 Parity and spectral flow

The spectral flow is a more refined invariant for paths of selfadjoint operators. For $x \in H^s(\mathbb{R}/\mathbb{Z})$ we use the Fourier expansion $x = \sum_{k \in \mathbb{Z}} e^{2\pi Jkt} x_k$ and $\sum_{k \in \mathbb{Z}} |k|^{2s} |x_k|^2 < \infty$. From the functional calculus of the selfadjoint operator

$$-J \frac{d}{dt} x = \sum_{k \in \mathbb{Z}} (2\pi k) e^{2\pi Jkt} x_k,$$

we define the selfadjoint operators

$$N_\mu x = \sum_{k \in \mathbb{Z}} (2\pi |k| + \mu) e^{2\pi Jkt} x_k, \quad \text{and} \quad P_\mu x = \sum_{k \in \mathbb{Z}} \frac{2\pi k + \mu}{2\pi |k| + \mu} e^{2\pi Jkt} x_k. \quad (3.6)$$

For $\mu > 0$ and $\mu \neq 2\pi k$, $k \in \mathbb{Z}$, the operator P_μ is an isomorphism on $H^s(\mathbb{R}/\mathbb{Z})$, for all $s \geq 0$.⁴ Consider the path

$$C(\eta) = P_\mu A(\eta) = P_\mu - N_\mu^{-1}(S(t; \eta) + \mu), \quad (3.7)$$

which is a path of operators of Fredholm index 0. The constant path $\eta \mapsto M_\mu(\eta) = P_\mu^{-1}$ is a parametrix for $\eta \mapsto C(\eta)$ (see [5,6]) and since $M_\mu C(\eta) = A(\eta)$, the parity of $C(\eta)$ is given by

$$\text{parity}(C(\eta), I) = \text{parity}(A(\eta), I). \quad (3.8)$$

Using N_μ , with $\mu > 0$ and $\mu \neq 2\pi k$, we define an equivalent norm on the Sobolev spaces $H^s(\mathbb{R}/\mathbb{Z})$:

$$(x, y)_{H^s} := (N_\mu^s x, N_\mu^s y)_{L^2}, \quad \forall x, y \in H^s(\mathbb{R}/\mathbb{Z}).$$

Lemma 3.4. *The operators $C(\eta)$ are selfadjoint on $(H^{1/2}(\mathbb{R}/\mathbb{Z}), (\cdot, \cdot)_{H^{1/2}})$ for all $\eta \in I$, and $\eta \mapsto C(\eta)$ is a path of selfadjoint operators on $H^{1/2}(\mathbb{R}/\mathbb{Z})$.*

Proof. From the functional calculus we derive that

$$(P_\mu x, y)_{H^s} = \sum_{k \in \mathbb{Z}} p_\mu(k) n_\mu^{2s}(k) x_k y_k = (x, P_\mu y)_{H^s},$$

where $n_\mu(k) = 2\pi|k| + \mu$ and $p_\mu(k) = \frac{2\pi k + \mu}{2\pi|k| + \mu}$. For $s = 1/2$ we have that

$$\begin{aligned} (N_\mu^{-1}(S(t; \eta) + \mu)x, y)_{H^{1/2}} &= ((S(t; \eta) + \mu)x, y)_{L^2} = (x, (S(t; \eta) + \mu)y)_{L^2} \\ &= (x, N_\mu^{-1}(S(t; \eta) + \mu)y)_{H^{1/2}}, \end{aligned}$$

which completes the proof. $\square$

For a path $\eta \mapsto \Lambda(\eta)$ of *selfadjoint* operators on a Hilbert space $\mathcal{H}$, which is continuously differentiable in the (strong) operator topology we define the crossing operator $\Gamma(\Lambda, \eta) = \pi \frac{d}{d\eta} \Lambda(\eta) \pi|_{\ker \Lambda(\eta)}$, where π is the orthogonal projection onto $\ker \Lambda(\eta)$. A crossing $\eta_0 \in I$ is a number for which the operator $\Lambda(\eta_0)$ is not invertible. A crossing is regular if $\Gamma(\Lambda, \eta_0)$ is non-singular. A point η_0 for which $\dim \ker \Lambda(\eta_0) = 1$, is called a simple crossing. A path $\eta \mapsto \lambda(\eta)$ is called generic if all crossings are simple. A path $\eta \mapsto \Lambda(\eta)$ with invertible endpoints can always be chosen to be generic by a small perturbation. At a simple crossing η_0 , there exists a C^1 -curve $\lambda(\eta)$, for η near η_0 , and $\lambda(\eta)$ is an eigenvalue of $\Lambda(\eta)$, with $\lambda(\eta_0) = 0$ and $\lambda'(\eta_0) \neq 0$, see [10,11]. The spectral flow for a generic path is defined by

$$\text{specflow}(\Lambda(\eta), I) = \sum_{\lambda(\eta_0)=0} \text{sgn}(\lambda'(\eta_0)). \quad (3.9)$$

⁴As before $\|P_\mu x\|_{H^s} \leq \|x\|_{H^s}$ and $\|P_\mu^{-1}x\|_{H^{1/2}} \leq C(\mu)\|x\|_{H^{1/2}}$, $\mu > 0$ and $\mu \neq 2\pi k$.

For a simple crossing η_0 the crossing operator is simply multiplication by $\lambda'(\eta_0)$ and

$$\Gamma(\Lambda, \eta)\psi(\eta_0) = \left(\frac{d}{d\eta} \Lambda(\eta_0)\psi(\eta_0), \psi(\eta_0) \right)_{\mathcal{H}} \psi(\eta_0) = \lambda'(\eta_0)\psi(\eta_0), \quad (3.10)$$

where $\psi(\eta_0)$ is normalized in $\mathcal{H}$, and

$$\lambda'(\eta_0) = \left(\frac{d}{d\eta} \Lambda(\eta_0)\psi(\eta_0)\psi(\eta_0) \right)_{\mathcal{H}}. \quad (3.11)$$

The spectral flow is defined for any continuously differentiable path $\eta \mapsto \Lambda(\eta)$ with invertible endpoints. From the theory in [6] there is a connection between the spectral flow of $\Lambda(\eta)$ and its parity:

$$\text{parity}(\Lambda(\eta), I) = (-1)^{\text{specflow}(\Lambda(\eta), I)}, \quad (3.12)$$

which in view of Equation (3.3) follows from the fact that $\text{cross}(\Lambda(\eta), I) = \text{specflow}(\Lambda(\eta), \eta) \bmod 2$ in the generic case.

The path $\eta \mapsto C(\eta)$ defined in (3.7) is a continuously differentiable path of operators on $H = H^{1/2}(\mathbb{R}/\mathbb{Z})$ with invertible endpoints, and therefore both parity and spectral flow are well-defined. If we combine Equations (3.5) and (3.8) with Equation (3.12) we obtain

$$\deg_{LS}(\Phi_{\mu, H}, B_\epsilon(x), 0) = \text{parity}(A(\eta), I) = (-1)^{\text{specflow}(C(\eta), I)}. \quad (3.13)$$

In the next section we link the spectral flow of $C(\eta)$ to the Conley-Zehnder indices of non-degenerate zeroes and therefore to the Euler-Floer characteristic.

4 The Conley-Zehnder index

We discuss the Conley-Zehnder index for Hamiltonian systems and mechanical systems, and explain the relation with the local degree and the Morse index for mechanical systems.

4.1 Hamiltonian systems

For a non-degenerate 1-periodic solution $x(t)$ of the Hamilton equations the Conley-Zehnder index can be defined as follows. The linearized flow Ψ is given by

$$\begin{cases} -J \frac{d\Psi}{dt} - D_x^2 H(x, t)\Psi = 0 \\ \Psi(0) = \text{Id}, \end{cases}$$

By Lemma 3.1(iii), a 1-periodic solution is non-degenerate if $\Psi(1)$ has no eigenvalues equal to 1. The Conley-Zehnder index is defined using the symplectic path $\Psi(t)$. Following [11], consider the crossing form $\Gamma(\Psi, t)$, defined for vectors $\xi \in \ker(\Psi(t) - \text{Id})$,

$$\Gamma(\Psi, t)\xi = \omega\left(\xi, \frac{d}{dt}\Psi(t)\xi\right) = (\xi, D_x^2 H(x(t), t)\xi). \quad (4.1)$$

A crossing $t_0 > 0$ is defined by $\det(\Psi(t_0) - \text{Id}) = 0$. A crossing is regular if the crossing form is non-singular. A path $t \mapsto \Psi(t)$ is regular if all crossings are regular. Any path can be approximated by a regular path with the same endpoints and which is homotopic to the initial path, see [10] for details. For a regular path $t \mapsto \Psi(t)$ the Conley-Zehnder index is given by

$$\mu^{CZ}(\Psi) = \frac{1}{2} \text{sgn } D_x^2 H(x(0), 0) + \sum_{\substack{t_0 > 0, \\ \det(\Psi(t_0) - \text{Id}) = 0}} \text{sgn } \Gamma(\Psi, t_0). \quad (4.2)$$

For a non-degenerate 1-periodic solution $x(t)$ we define the Conley-Zehnder index as $\mu^{CZ}(x) := \mu^{CZ}(\Psi)$, and the index is integer valued.

Let x be a 1-periodic solution and consider the path $\eta \mapsto B(\eta; x) = -J \frac{d}{dt} - S(t; \eta)$, where, as before, $S(t; \eta)$ is a smooth path of symmetric matrices with endpoints $S(t; 0) = \theta \text{Id}$ and $S(t; 1) = D_x^2 H(x(t), t)$ with $\theta \neq 2\pi k, k \in \mathbb{Z}$. The operators $B(\eta) = B(\eta; x)$ are unbounded operators on $L^2(\mathbb{R}/\mathbb{Z})$, with domain $H^1(\mathbb{R}/\mathbb{Z})$. A path $\eta \mapsto B(\eta)$ is continuously differentiable in the (weak) operator topology of $\mathcal{B}(H^1, L^2)$ and Hypotheses (A1)-(A3) in [11] are satisfied. We now repeat the definition of spectral flow for a path of unbounded operators as developed in [11]. The crossing operator for a path $\eta \mapsto B(\eta)$ is given by $\Gamma(B, \eta) = \pi \frac{d}{d\eta} B(\eta) \pi|_{\ker B(\eta)}$, where π is the orthogonal projection onto $\ker B(\eta)$. A crossing $\eta_0 \in I$ is a number for which the operator $B(\eta_0)$ is not invertible. A crossing is regular if $\Gamma(B, \eta_0)$ is non-singular. A point η_0 for which $\dim \ker B(\eta_0) = 1$, is called a simple crossing. A path $\eta \mapsto B(\eta)$ is called generic if all crossing are simple. A path $\eta \mapsto B(\eta)$ can always be chosen to be generic. At a simple crossing η_0 there exists a C^1 -curve $\ell(\eta)$, for η near η_0 , and $\ell(\eta)$ is an eigenvalue of $B(\eta)$ with $\ell(\eta_0) = 0$ and $\ell'(\eta_0) \neq 0$. The spectral flow for a generic path is defined by

$$\text{specflow}(B(\eta), I) = \sum_{\ell(\eta_0)=0} \text{sgn}(\ell'(\eta_0)), \quad (4.3)$$

and at simple crossings η_0 ,

$$\Gamma(B, \eta)\phi(\eta_0) = \left(\frac{d}{d\eta} B(\eta_0)\phi(\eta_0), \phi(\eta_0) \right)_{L^2} \phi(\eta_0) = \ell'(\eta_0)\phi(\eta_0), \quad (4.4)$$

after normalizing $\phi(\eta_0)$ in $L^2(\mathbb{R}/\mathbb{Z})$. As before the derivative of ℓ at η_0 is given by

$$\ell'(\eta_0) = -(\partial_\eta S(t; \eta_0)\phi(\eta_0), \phi(\eta_0))_{L^2}. \quad (4.5)$$

Proposition 4.1. Let $\eta \mapsto B(\eta), \eta \in I$, as defined above, be a generic path of unbounded self-adjoint operators with invertible endpoints, and let $\eta \mapsto \Psi(\eta; t)$ be the associated path of symplectic matrices defined by

$$\begin{cases} -J \frac{d\Psi}{dt}(t; \eta) - S(t; \eta) \Psi(t; \eta) = 0 \\ \Psi(0; \eta) = \text{Id}, \end{cases}$$

Then

$$\text{specflow}(B(\eta), I) = \mu_{B(0)}^{CZ} - \mu_{B(1)}^{CZ} \quad (4.6)$$

where $\mu_{B(0)}^{CZ} = \mu^{CZ}(\Psi(t; 0))$, $\mu_{B(1)}^{CZ} = \mu^{CZ}(\Psi(t; 1))$.

Proof. The expression for the spectral flow follows from [11] and [12]. $\square$

In the case $\eta = 0$, the Conley-Zehnder index $\mu_{B(0)}^{CZ}$ can be computed explicitly. Recall that $B(0) = -J \frac{d}{dt} - S(0) = -J \frac{d}{dt} - \theta \text{Id}$.

Lemma 4.1. Let $\theta > 0$ (fixed) and $\theta \neq 2\pi k$, then $\mu_{B(0)}^{CZ} = 1 + 2 \lfloor \frac{\theta}{2\pi} \rfloor$.

Proof. The solution to $B(0)\Psi(t) = 0$ is given by $\Psi(t) = e^{\theta J t}$ and $\det(\Psi(1) - \text{Id}) = 0$ exactly when $t = t_0 = \frac{2\pi k}{\theta}$. By (4.1) and (4.2) we have that $\Gamma(\Psi, t)\xi = \theta|\xi|^2$ and therefore $\mu_{B(0)}^{CZ} = 1 + 2 \lfloor \frac{\theta}{2\pi} \rfloor$, which proves the lemma. $\square$

The zeroes $x \in \Phi_{\mu, H}^{-1}(0)$ in $\Omega = [x] \text{rel } y$ can be estimated by Braid Floer homology $\text{HB}_*([x] \text{rel } y)$ of $\Omega = [x] \text{rel } y$. The Euler-Floer characteristic of $\text{HB}_*([x] \text{rel } y)$ is defined as

$$\chi(\text{HB}_*([x] \text{rel } y)) := \sum_{k \in \mathbb{Z}} (-1)^k \dim \text{HB}_k([x] \text{rel } y). \quad (4.7)$$

In [12] the following analogue of the Poincaré-Hopf formula is proved.

Proposition 4.2. For a proper braid class $[x] \text{rel } y$ and a generic Hamiltonian $H \in \mathcal{H}_{\parallel}^{\text{reg}}(y)$, it holds that

$$\chi(\text{HB}_*([x] \text{rel } y)) = \sum_{x \in \Phi_{\mu, H}^{-1}(0)} (-1)^{\mu^{CZ}(x)}.$$

It remains to show that $\chi(\text{HB}_*([x] \text{rel } y))$ and $\deg_{LS}(\Phi_{\mu, H}, \Omega, 0)$ are related.

Proposition 4.3. For a proper braid class $[x] \text{rel } y$ and a generic Hamiltonian $H \in \mathcal{H}_{\parallel}^{\text{reg}}(y)$, we have that

$$\chi(\text{HB}_*([x] \text{rel } y)) = - \sum_{x_i \in \Phi_{\mu, H}^{-1}(0)} (-1)^{-\text{specflow}(B(\eta; x), I)}, \quad (4.8)$$

where $\eta \mapsto B(\eta; x)$ is given above for $x \in \Phi_{\mu, H}^{-1}(0)$.

Proof. By Proposition 4.1 and Lemma 4.1 the spectral flow satisfies,

$$\begin{aligned}\mu^{CZ}(x) &= \mu_{B(1;x)}^{CZ} = \mu_{B(0)}^{CZ} - \text{specflow}(B(\eta; x), I) \\ &= 1 + 2\left\lfloor \frac{\theta}{2\pi} \right\rfloor - \text{specflow}(B(\eta; x), I).\end{aligned}$$

This implies

$$(-1)^{\mu^{CZ}(x)} = -(-1)^{-\text{specflow}(B(\eta; x), I)},$$

which completes the proof. $\square$

4.2 Mechanical systems

A mechanical system is defined as the Euler-Lagrange equations of the Lagrangian density $L(q, t) = \frac{1}{2}q_t^2 - V(q, t)$. The linearization at a critical points $q(t)$ of the Lagrangian action is given by the unbounded operator

$$-\frac{d^2}{dt^2} - D_q^2 V(q(t), t) : H^2(\mathbb{R}/\mathbb{Z}) \subset L^2(\mathbb{R}/\mathbb{Z}) \rightarrow L^2(\mathbb{R}/\mathbb{Z}).$$

Consider a path of unbounded self-adjoint operators on $L^2(\mathbb{R}/\mathbb{Z})$ given by $\eta \mapsto D(\eta) = -\frac{d^2}{dt^2} - Q(t; \eta)$, with $Q(t; \eta)$ smooth. If $D(0)$ and $D(1)$ are invertible, then the spectral flow is well-defined.

Proposition 4.4. *Assume that the endpoints of $\eta \mapsto D(\eta)$ are invertible. Then*

$$\text{specflow}(D(\eta), I) = \beta_{D(0)} - \beta_{D(1)}, \quad (4.9)$$

where $\beta_{D(0)}$ and $\beta_{D(1)}$ are the Morse indices of $D(0)$ and $D(1)$ respectively.

Proof. In [11] the concatenation property of the spectral flow is proved. We use concatenation as follows. Let $c > 0$ be a sufficiently large constant such that $D(0) + c \text{Id}$ and $D(1) + c \text{Id}$ are positive definite self-adjoint operators on $L^2(\mathbb{R}/\mathbb{Z})$. Consider the paths $\eta \mapsto D_1(\eta) = D(0) + \eta c \text{Id}$ and $\eta \mapsto D_2(\eta) = D(1) + (1 - \eta)c \text{Id}$. Their concatenation $D_1 \# D_2$ is a path from $D(0)$ to $D(1)$ and $\eta \mapsto D_1 \# D_2$ is homotopic to $\eta \mapsto D(\eta)$. Using the homotopy invariance and the concatenation property of the spectral flow we obtain

$$\text{specflow}(D(\eta), I) = \text{specflow}(D_1 \# D_2, I) = \text{specflow}(D_1, I) + \text{specflow}(D_2, I).$$

Since $D(0)$ is invertible, the regular crossings of $D_1(\eta)$ are given by $\eta_i^1 = -\frac{\lambda_i}{c}$, where λ_i are negative eigenvalues of $D(0)$. By the positive definiteness of $D(0) + c \text{Id}$, the negative eigenvalues of $D(0)$ satisfy $0 > \lambda_i > -c$. For the crossing η_i this implies

$$0 < \eta_i = -\frac{\lambda_i}{c} < 1,$$

and therefore the number of crossings equals the number of negative eigenvalues of $D(0)$ counted with multiplicity. By the choice of c , we also have that $\frac{d}{d\eta}D_1(\eta) = c\text{Id}$ is positive definite and therefore the signature of the crossing operator of $D_1(\eta)$ is exactly the number of negative eigenvalues of $D(0)$, i.e. $\text{specflow}(D_1, I) = \beta_{D(0)}$. For $D_2(\eta)$ we obtain, $\text{specflow}(D_2, I) = -\beta_{D(1)}$. This proves that $\text{specflow}(D(\eta), I) = \beta_{D(0)} - \beta_{D(1)}$. $\square$

For a mechanical system we have the Hamiltonian $H(x, t) = \frac{1}{2}p^2 + V(q, t)$. As such the Conley-Zehnder index of a critical point q can be defined as the Conley-Zehnder index of $x = (q_t, q)$ using the mechanical Hamiltonian, see also [1] and [2].

Lemma 4.2. *Let q be a critical point of the mechanical Lagrangian action, then the associated Conley-Zehnder index $\mu^{CZ}(x)$ is well-defined, and $\mu^{CZ}(x) = \beta(q)$, where $\beta(q)$ is the Morse index of q .*

Proof. As before, consider the curves $\eta \mapsto B(\eta)$ and $\eta \mapsto D(\eta)$, $\eta \in I = [0, 1]$ given by

$$B(\eta) = -J \frac{d}{dt} - \begin{pmatrix} 1 & 0 \\ 0 & Q(t; \eta) \end{pmatrix}, \quad D(\eta) = -\frac{d^2}{dt^2} - Q(t; \eta).$$

The crossing forms of the curves are the same — $\Gamma(B, \eta) = \Gamma(D, \eta)$ — and therefore also the crossings η_0 are identical. Indeed, $B(\eta_0)$ is non-invertible if and only if $D(\eta_0)$ is non-invertible. Consequently, $\text{specflow}(B(\eta), I) = \text{specflow}(D(\eta), I)$ and the Propositions 4.1 and 4.4 then imply that

$$\beta_{D(0)} - \beta_{D(1)} = \mu_{B(0)}^{CZ} - \mu_{B(1)}^{CZ}. \quad (4.10)$$

Now choose $Q(t; \eta)$ such that $Q(t; 0) = d^2V(q(t), t) + c$ and $Q(t; 1) = D_q^2V(q(t), t)$ and such that $\eta \mapsto B(\eta)$ and $\eta \mapsto D(\eta)$ are regular curves. If $c \ll 0$, then $\beta_{D(0)} = 0$. In order to compute $\mu_{B(0)}^{CZ}$ we invoke the crossing from $\Gamma(\Psi, t)$ for the associated symplectic path $\Psi(t)$ as explained in Section 4. Crossings at $t_0 \in (0, 1]$ correspond to non-trivial solutions of the equation $D(0)\psi = 0$ on $[0, t_0]$, with periodic boundary conditions. To be more precise, let $\Psi = (\phi, \psi)$, then $B(0)\Psi = 0$ is equivalent to $\psi_t = \phi$ and $-\phi_t - (D_q^2V(q(t), t) + c)\psi = 0$, which yields the equation $D(0)\psi = 0$. For the latter the kernel is trivial for any $t_0 \in (0, 1]$. Indeed, if ψ is a solution, then $\int_0^{t_0} |\psi_t|^2 = \int_0^{t_0} (D_q^2V(q, t) + c)\psi^2 < 0$, which is a contradiction. Therefore, there are no crossing $t_0 \in (0, 1]$. As for $t_0 = 0$ we have that $(D_q^2V(q(0), 0) + c) < 0$, which implies that $\text{sgn } S(0; 0) = 0$ and therefore $\mu_{B(0)}^{CZ} = 0$, which proves the lemma. $\square$

5 The spectral flows are the same

In order to show that the spectral flows are the same we use the fact that the paths $\eta \mapsto C(\eta)$ and $\eta \mapsto B(\eta)$ for a non-degenerate zero $x_i \in \Phi_{\mu, H}^{-1}(0)$ are chosen to have only simple crossings for their crossing operators, i.e. zero eigenvalues are simple. In this case the spectral flows are determined by the signs of the derivatives of the eigenvalues at the crossings. For $\eta \mapsto B(\eta)$ the expression given by Equation (4.5) and from Equation (3.11) a similar expression for $\eta \mapsto C(\eta)$ can be derived and is given by

$$\lambda'(\eta_0) = -(N_\mu^{-1} \partial_\eta S(t; \eta_0) \psi(\eta_0), \psi(\eta_0))_{H^{1/2}} = -(\partial_\eta S(t; \eta_0) \psi(\eta_0), \psi(\eta_0))_{L^2} \quad (5.1)$$

Lemma 5.1. *The sets $\{\eta \in [0, 1] : C(\eta) \psi(\eta) = 0\}$ and $\{\eta \in [0, 1] : B(\eta) \phi(\eta) = 0\}$ are the same, and the operators $C(\eta)$ and $B(\eta)$ have the same eigenfunctions at crossings η_0 . In particular, $\eta \mapsto B(\eta)$ is generic if and only if $\eta \mapsto C(\eta)$ is generic.*

Proof. Given $\eta_0 \in [0, 1]$ such that $C(\eta_0) \psi(\eta_0) = 0$, then

$$P_\mu \psi(\eta_0) - N_\mu^{-1} (S(\eta_0; t) + \mu) \psi(\eta_0) = 0,$$

and thus $\psi(\eta_0) - L_\mu^{-1} (S(\eta_0; t) + \mu) \psi(\eta_0) = 0$, which is equivalent to the equation $(-J \frac{d}{dt} - S(t; \eta_0)) \psi(\eta_0) = 0$, i.e. $B(\eta_0) \psi(\eta_0) = 0$. $\square$

Lemma 5.2. *For all $\mu > 0$, with $\mu \neq 2\pi k$, $k \in \mathbb{Z}$, $\text{sgn } \lambda'(\eta_0) = \text{sgn } \ell'(\eta_0)$ for all crossings at η_0 .*

Proof. The eigenfunctions $\psi(\eta_0)$ in Equation (5.1) for $\lambda'(\eta_0)$ are normalized in $H^{1/2}(\mathbb{R}/\mathbb{Z})$ and therefore they relate to the eigenfunctions $\phi(\eta_0)$ in Equation (4.5) for $\ell'(\eta_0)$ as follows:

$$\psi(\eta_0) = \frac{\phi(\eta_0)}{\|\phi(\eta_0)\|_{H^{1/2}}}, \quad \|\phi(\eta_0)\|_{L^2} = 1.$$

Combining Equations (4.5) and (5.1) then gives

$$\begin{aligned} \lambda'(\eta_0) &= -(\partial_\eta S(t; \eta_0) \psi(\eta_0), \psi(\eta_0))_{L^2} \\ &= -\frac{1}{\|\phi(\eta_0)\|_{H^{1/2}}^2} (\partial_\eta S(t; \eta_0) \phi(\eta_0), \phi(\eta_0))_{L^2} = \frac{\ell'(\eta_0)}{\|\phi(\eta_0)\|_{H^{1/2}}^2}, \end{aligned}$$

which proves the lemma. $\square$

Lemma 5.2 implies that for any non-degenerate $x \in \Phi_{\mu, H}^{-1}(0) \cap \Omega$

$$\text{specflow}(C(\eta; x), I) = \text{specflow}(B(\eta; x), I), \quad (5.2)$$

where $B(\eta; x)$ and $C(\eta; x)$ are the above described path associated with x . Therefore

$$\text{parity}(A(\eta; x), I) = (-1)^{\text{specflow}(C(\eta; x), I)} = (-1)^{\text{specflow}(B(\eta; x), I)}, \quad (5.3)$$

which yields the following proposition.

Proposition 5.1. *The Leray-Schauder degree satisfies*

$$\deg_{LS}(\Phi_{\mu, H}, \Omega, 0) = -\chi(\text{HB}_*([x] \text{ rel } y)).$$

Proof. For any Hamiltonian $H \in \mathcal{H}_{\parallel}(y)$ there exists a generic Hamiltonian $\tilde{H} \in \mathcal{H}_{\parallel}^{\text{reg}}(y)$ such all zeroes $x_i \in \Phi_{\mu, \tilde{H}}^{-1}(0) \cap \Omega$ are non-degenerate. Since $\Omega = [x] \text{ rel } y$ is isolating for all Hamiltonians in $\mathcal{H}_{\parallel}(y)$, the invariance of the Leray-Schauder degree yields $\deg_{LS}(\Phi_{\mu, H}, \Omega, 0) = \deg_{LS}(\Phi_{\mu, \tilde{H}}, \Omega, 0)$. From the Propositions 3.3 and 4.3 and Equation (5.3) we conclude that

$$\begin{aligned} \deg_{LS}(\Phi_{\mu, H}, \Omega, 0) &= \deg_{LS}(\Phi_{\mu, \tilde{H}}, \Omega, 0) \\ &= \sum_{x \in \Phi_{\mu, \tilde{H}}^{-1}(0)} \deg_{LS}(\Phi_{\mu, \tilde{H}}, B_{\epsilon}(x), 0) = \sum_{x \in \Phi_{\mu, \tilde{H}}^{-1}(0)} \text{parity}(A(\eta; x), I) \\ &= \sum_{x \in \Phi_{\mu, \tilde{H}}^{-1}(0)} (-1)^{\text{specflow}(B(\eta; x), I)} = \sum_{x \in \Phi_{\mu, \tilde{H}}^{-1}(0)} (-1)^{-\text{specflow}(B(\eta; x), I)} \\ &= -\chi(\text{HB}_*([x] \text{ rel } y)), \end{aligned}$$

which completes the proof. $\square$

Remark 5.1. As $\mu \gg 1$ it holds that $\ell'(\eta_0) \sim \mu \lambda'(\eta_0)$. Indeed, $\|\phi(\eta_0)\|_{H^{1/2}}^2 = \sum_k (2\pi|k| + \mu) a_k^2$, where a_k are the Fourier coefficients of $\phi(\eta_0)$ and $\sum_k a_k^2 = 1$. Since $\phi(\eta_0)$ are smooth functions the Fourier coefficients satisfy the following properties. For any $\delta > 0$ and any $s > 0$, there exists $N_{\delta, s} > 0$ such that $\sum_{|k| \geq N} |k|^{2s} |a_k|^2 \leq \delta$, for all $N \geq N_{\delta, s}$. From the latter it follows that $\sum_k 2\pi|k| a_k^2 \leq C$, with $C > 0$ independent of η_0 and μ . We derive that $\mu \leq \|\phi(\eta_0)\|_{H^{1/2}}^2 \leq C + \mu$ and

$$1 \leftarrow \frac{\mu}{C + \mu} \leq \frac{\mu \lambda'(\eta_0)}{\ell'(\eta_0)} = \frac{\mu}{\|\phi(\eta_0)\|_{H^{1/2}}^2} \leq \frac{\mu}{\mu} = 1,$$

as $\mu \rightarrow \infty$, which proves our statement.

6 The proof of Theorems 1.1 and 1.2

We start with the proof of Theorem 1.2. Since $\text{HB}_*([x] \text{ rel } y)$ is an invariant of the proper braid class $[x \text{ rel } y]$ it does not depend on a particular fiber $[x] \text{ rel } y$. Therefore

we denote the Euler-Floer characteristic by $\chi(x \text{ rel } y) := \chi(\text{HB}_*([x] \text{ rel } y))$. Recall the homotopy invariance of the Leray-Schauder degree as expressed in Equation (2.3)

$$\deg_{LS}(\Phi_\mu, \Omega, 0) = \deg_{LS}(\Phi_{\mu, \alpha}, \Omega, 0) = \deg_{LS}(\Phi_{\mu, H}, \Omega, 0).$$

By Proposition 5.1 we have that

$$\deg_{LS}(\Phi_\mu, \Omega, 0) = \deg_{LS}(\Phi_{\mu, H}, \Omega, 0) = -\chi(x \text{ rel } y),$$

and $\chi(x \text{ rel } y) \neq 0$, then implies that $\Phi_\mu^{-1}(0) \cap \Omega \neq \emptyset$. Therefore there exists a closed integral curves in any relative braid class fiber of $[x \text{ rel } y]$, whenever $\chi(x \text{ rel } y) \neq 0$, and this completes the proof of Theorem 1.2.

The remainder of this section is to prove the Poincaré-Hopf Formula in Theorem 1.1 for closed integral curves in proper braid fibers. The mapping

$$\mathcal{E} : C^1(\mathbb{R}/\mathbb{Z}) \rightarrow C^0(\mathbb{R}/\mathbb{Z}), \quad \mathcal{E}(x) = \frac{dx}{dt} - X(x, t),$$

is smooth (nonlinear) Fredholm mapping of index 0. Let $M \in \text{GL}(C^0, C^1)$ be an isomorphism such that $M\mathcal{E}(x)$ is of the form $M\mathcal{E}(x) = \Phi_M(x) = x - K_M(x)$, with $K_M : C^1(\mathbb{R}/\mathbb{Z}) \rightarrow C^1(\mathbb{R}/\mathbb{Z})$ compact. Such isomorphisms M (constant parametrices) obviously exist. For example $M = \left(\frac{d}{dt} + 1\right)^{-1}$, or $M = -JL_\mu^{-1}$. The mappings $\Phi_M : C^1(\mathbb{R}/\mathbb{Z}) \rightarrow C^1(\mathbb{R}/\mathbb{Z})$ are Fredholm mappings of index 0.

Let $x \in C^1(\mathbb{R}/\mathbb{Z})$ be a non-degenerate zero of $\mathcal{E}$ and recall the index $\iota(x)$:

$$\iota(x) = -\text{sgn}(\det(\Theta))(-1)^{\beta_M(\Theta)} \deg_{LS}(\Phi_M, B_\epsilon(x), 0),$$

where $\Theta \in M_{2 \times 2}(\mathbb{R})$, with $\sigma(\Theta) \cap 2\pi k i \mathbb{R} = \emptyset$, $k \in \mathbb{Z}$ and $\beta_M(\Theta)$ is the Morse index of $\text{Id} - K_M(0)$.

Lemma 6.1. *The index $\iota(x)$ for a non-degenerate zero of $\mathcal{E}$ is well-defined, i.e. independent of the choices of $M \in \text{GL}(C^0, C^1)$ and $\Theta \in M_{2 \times 2}(\mathbb{R})$.*

Proof. Consider smooth paths $\eta \mapsto F_\Theta(\eta)$, defined by $F_\Theta(\eta) = \frac{d}{dt} - R(t; \eta)$, where $R(t; 0) = \Theta$ and $R(t; 1) = D_x X(x(t), t)$. The path

$$F_\Theta : [0, 1] \rightarrow \text{Fred}_0(C^1, C^0)$$

has invertible end points, and by the theory in [3, 4] we have that the parity of $\eta \mapsto F_\Theta(\eta)$ is well-defined and independent of M , i.e.

$$\begin{aligned} \text{parity}(F_\Theta(\eta), I) &= \text{parity}(D_{M, \Theta}(\eta), I) = (-1)^{\beta_M(\Theta)} (-1)^{\beta_M(x)} \\ &= (-1)^{\beta_M(\Theta)} \deg_{LS}(\Phi_M, B_\epsilon(x), 0), \end{aligned}$$

where $D_{M,\Theta}(\eta) = MF_{\Theta}(\eta)$ and $\beta_M(x)$ is the Morse index of $D_{M,\Theta}(1) = \text{Id} - K_M(1)$. It remains to show that the index $\iota(x)$ is independent with respect to Θ . Let Θ and Θ' be admissible matrices and let $\eta \mapsto G(\eta)$ be a path connecting $G(0) = \frac{d}{dt} - \Theta$ and $G(1) = \frac{d}{dt} - \Theta'$. For the parities it holds that

$$\text{parity}(F_{\Theta}(\eta), I) = \text{parity}(G(\eta), I) \cdot \text{parity}(F_{\Theta'}(\eta), I).$$

To compute $\text{parity}(G(\eta), I)$ we consider a special parametrix M_{μ} , given by $M_{\mu} = \left(\frac{d}{dt} + \mu\right)^{-1}$, $\mu > 0$. From the definition of parity we have that

$$\text{parity}(G(\eta), I) = \text{parity}(M_{\mu}G(\eta), I) = \deg_{LS}(M_{\mu}G(0)) \cdot \deg_{LS}(M_{\mu}G(1)).$$

We now compute the Leray-Schauder degrees of $M_{\mu}G(0)$ and $M_{\mu}G(1)$. We start with Θ and in order to compute the degree we determine the Morse index. Consider the eigenvalue problem

$$M_{\mu}G(0)\psi = \lambda\psi, \quad \lambda \in \mathbb{R},$$

which is equivalent to $(1 - \lambda)\frac{d\psi}{dt} = (\Theta + \lambda\mu)\psi$. Non-trivial solutions are given by $\psi(t) = \exp\left(\frac{\Theta + \lambda\mu}{1 - \lambda}t\right)\psi_0$, which yields the condition $\frac{\theta + \lambda\mu}{1 - \lambda} = 2\pi ki$, $k \in \mathbb{Z}$, where θ is an eigenvalue of Θ . We now consider three cases:

(i) $\theta_{\pm} = a \pm ib$. In case of a negative eigenvalue λ we have $\frac{a + \lambda\mu}{1 - \lambda} = 0$ and $\frac{b}{1 - \lambda} = 2\pi k$. The same $\lambda < 0$ also suffices for the conjugate eigenvalue via $\frac{-b}{1 - \lambda} = -2\pi k$. This implies that any eigenvalue $\lambda < 0$ has multiplicity 2, and thus $\deg_{LS}(M_{\mu}G(0)) = 1$.

(ii) $\theta_{\pm} \in \mathbb{R}$, $\theta_- \cdot \theta_+ > 0$. In case of a negative eigenvalue λ we have $\frac{\theta_{\pm} + \lambda\mu}{1 - \lambda} = 0$ and thus $\lambda_{\pm} = -\frac{\theta_{\pm}}{\mu}$, which yields two negative or two positive eigenvalues. As before $\deg_{LS}(M_{\mu}G(0)) = 1$.

(iii) $\theta_{\pm} \in \mathbb{R}$, $\theta_- \cdot \theta_+ < 0$. From case (ii) we easily derive that there exist two eigenvalues $\lambda_{\pm}$, one positive and one negative, and therefore $\deg_{LS}(M_{\mu}G(0)) = -1$.

These cases combined imply that $\deg_{LS}(M_{\mu}G(0)) = \text{sgn}(\det(\Theta))$ and

$$\text{parity}(G(\eta), I) = \text{sgn}(\det(\Theta)) \cdot \text{sgn}(\det(\Theta')).$$

From the latter we derive:

$$\begin{aligned} & \text{sgn}(\det(\Theta)) \cdot \text{parity}(F_{\Theta}(\eta), I) \\ &= \text{sgn}(\det(\Theta)) \cdot \text{sgn}(\det(\Theta)) \cdot \text{sgn}(\det(\Theta')) \cdot \text{parity}(F_{\Theta'}(\eta), I) \\ &= \text{sgn}(\det(\Theta')) \cdot \text{parity}(F_{\Theta'}(\eta), I), \end{aligned}$$

which proves the independence of Θ . □

Lemmas 6.1 shows that the index of a non-degenerate zero of $\mathcal{E}$ is well-defined. We now show that the same holds for isolated zeroes.

Lemma 6.2. *The index $\iota(x)$ for an isolated zero of $\mathcal{E}$ is well-defined and for a fixed choice of M and Θ the index is given by*

$$\iota(x) = -\operatorname{sgn}(\det(\Theta))(-1)^{\beta_M(\Theta)} \deg_{LS}(\Phi_M, B_\epsilon(x), 0),$$

where $\epsilon > 0$ is small enough such that x is the only zero of $\mathcal{E}$ in $B_\epsilon(x)$.

Proof. By the Sard-Smale Theorem one can choose an arbitrarily small $h \in C^0(\mathbb{R}/\mathbb{Z})$, $\|h\|_{C^0} < \epsilon'$, such that h is a regular value of $\mathcal{E}$ and $\mathcal{E}^{-1}(h) \cap B_\epsilon(x)$ consists of finitely many non-degenerate zeroes in x_h . Set $\tilde{\mathcal{E}}(x) = \mathcal{E}(x) - h$ and define

$$\iota(x) = \sum_{x_h \in \tilde{\mathcal{E}}^{-1}(0) \cap B_\epsilon(x)} \iota(x_h). \quad (6.1)$$

We now show that $\iota(x)$ is well-defined. Choose a fixed parametrix M (for $\mathcal{E}$) and fixed $\Theta \in M_{2 \times 2}(\mathbb{R})$, and let $\tilde{\Phi}_M = M\tilde{\mathcal{E}}$, then

$$\sum_{x_h} \iota(x_h) = -\operatorname{sgn}(\det(\Theta))(-1)^{\beta_M(\Theta)} \sum_{x_h} \deg_{LS}(\tilde{\Phi}_M, B_{\epsilon_h}(x_h), 0),$$

where $B_{\epsilon_h}(x_h)$ are sufficiently small neighborhoods containing only one zero. From Leray-Schauder degree theory we derive that

$$\sum_{x_h} \deg_{LS}(\tilde{\Phi}_M, B_{\epsilon_h}(x_h), 0) = \deg_{LS}(\tilde{\Phi}_M, B_\epsilon(x), 0) = \deg_{LS}(\Phi_M, B_\epsilon(x), 0),$$

which proves the lemma. $\square$

Theorem 1.1 now follows from the Leray-Schauder degree. Suppose all zeroes of $\mathcal{E}$ in $\Omega = [x] \operatorname{rel} y$ are isolated, then Lemma 6.2 implies that

$$\begin{aligned} \sum_{x \in \mathcal{E}^{-1}(0) \cap \Omega} \iota(x) &= -\operatorname{sgn}(\det(\Theta))(-1)^{\beta_M(\Theta)} \sum_x \deg_{LS}(\Phi_M, B_\epsilon(x), 0) \\ &= -\operatorname{sgn}(\det(\Theta))(-1)^{\beta_M(\Theta)} \deg_{LS}(\Phi_M, \Omega, 0) \end{aligned}$$

Since the latter expression is independent of M and Θ we choose $M = L_\mu^{-1}$ and $\Theta = \theta J$. Then, $\Phi_M = \Phi_\mu$, and for the indices we have $\operatorname{sgn}(\det(\theta J)) = 1$ and by Lemma 3.3, $(-1)^{\beta_{L_\mu^{-1}}(\theta J)} = 1$. By Proposition 5.1, $\deg_{LS}(\Phi_\mu, \Omega, 0) = -\chi(x \operatorname{rel} y)$, which, by substitution of these choices into the index formula, yields

$$\sum_{x \in \mathcal{E}^{-1}(0) \cap \Omega} \iota(x) = \chi(x \operatorname{rel} y),$$

completing the proof Theorem 1.1.

7 Computing the Euler-Floer characteristic

In section we prove Theorem 1.3 and show that the Euler-Floer characteristic can be determined via a discrete topological invariant.

7.1 Hyperbolic Hamiltonians on $\mathbb{R}^2$

Consider Hamiltonians of the form

$$H(x, t) = \frac{1}{2}p^2 - \frac{1}{2}q^2 + h(x, t), \quad (7.1)$$

where h satisfies the following hypotheses:

(h1) $h \in C^\infty(\mathbb{R}^2 \times \mathbb{R}/\mathbb{Z})$;

(h2) $\text{supp}(h) \subset \mathbb{R} \times [-R, R] \times \mathbb{R}/\mathbb{Z}$, for some $R > 0$;

(h3) $\|h\|_{C_b^2(\mathbb{R}^2 \times \mathbb{R}/\mathbb{Z})} \leq c$.

Lemma 7.1. *Let H be given by (7.1), with h satisfying (h1)-(h3). Then, there exists a constant $R' \geq R > 0$, such any 1-periodic solution of x of $x' = X_H(x, t)$ satisfies the estimate*

$$|x(t)| \leq R', \quad \text{for all } t \in \mathbb{R}/\mathbb{Z}.$$

Proof. The Hamilton equation in local coordinates are given by

$$p_t = q - h_q(p, q, t), \quad q_t = p + h_p(p, q, t).$$

Since h is smooth we can rewrite the equations as

$$q_{tt} = h_{pq}(p, q, t)q_t + (1 + h_{pp}(p, q, t))(q - h_q(p, q, t)) + h_{pt}(p, q, t). \quad (7.2)$$

If $x(t)$ is a 1-periodic solution to the Hamilton equations, and suppose there exists an interval $I = [t_0, t_1] \subset [0, 1]$ such that $|q(t)| > R$ on $\text{int}(I)$ and $|q(t)||_{\partial I} = R$. The function $q|_I$ satisfies the equation $q_{tt} - q = 0$, and obviously such solutions do not exist. Indeed, if $q|_I \geq R$, then $q_t(t_0) \geq 0$ and $q_t(t_1) \leq 0$ and thus $0 \geq q_t|_{\partial I} = \int_I q \geq R|I| > 0$, a contradiction. The same holds for $q|_I \leq -R$. We conclude that

$$|q(t)| < R, \quad \text{for all } t \in \mathbb{R}/\mathbb{Z}.$$

We now use the a priori q -estimate in combination with Equation (7.2) and Hypothesis (h3). Multiplying Equation (7.2) by q and integrating over $[0, 1]$ gives:

$$\begin{aligned} \int_0^1 q_t^2 &= - \int_0^1 h_{pq} q_t q - \int_0^1 (1 + h_{pp})(q - h_q) q - \int_0^1 h_{pt} q \\ &\leq C \int_0^1 |q_t| + C \leq \epsilon \int_0^1 q_t^2 + C_\epsilon, \end{aligned}$$

which implies that $\int_0^1 q_t^2 \leq C(R)$. The L^2 -norm of the right hand side in (7.2) can be estimated using the L^∞ estimate on q and the L^2 -estimate on q_t , which yields $\int_0^1 q_{tt}^2 \leq C(R)$. Combining these estimates we have that $\|q\|_{H^2(\mathbb{R}/\mathbb{Z})} \leq C(R)$ and thus $|q_t(t)| \leq C(R)$, for all $t \in \mathbb{R}/\mathbb{Z}$. From the Hamilton equations it follows that $|p(t)| \leq |q_t(t)| + C$, which proves the lemma. $\square$

Lemma 7.2. *If $H(x, t; \alpha)$, $\alpha \in [0, 1]$ is a (smooth) homotopy of Hamiltonians satisfying (h1)-(h3) with uniform constants $R > 0$ and $c > 0$, then $|x_\alpha(t)| \leq R'$, for all 1-periodic solutions and for all $\alpha \in [0, 1]$.*

Proof. The a priori H^2 -estimates in Lemma 7.1 hold with uniform constants with respect to $\alpha \in [0, 1]$. This then proves the lemma. $\square$

7.2 Braids on $\mathbb{R}^2$ and Legendrian braids

In Section 1 we defined braid classes as path components of closed loops in $\mathcal{LC}_n(\mathbb{D}^2)$, denoted by $[x]$. If we consider closed loops in $\mathbf{C}_n(\mathbb{R}^2)$, then the braid classes will be denoted by $[x]_{\mathbb{R}^2}$. The same notation applies to relative braid classes $[x \text{ rel } y]_{\mathbb{R}^2}$. A relative braid class is proper if components $x_c \subset x$ cannot be deformed onto (i) itself, or other components $x'_c \subset x$, or (ii) components $y_c \subset y$. A fiber $[x]_{\mathbb{R}^2} \text{ rel } y$ is *not* bounded!

In order to compute the Euler-Floer characteristic of $[x \text{ rel } y]$ we assume without loss of generality that $x \text{ rel } y$ is a positive representative. If not we compose $x \text{ rel } y$ with a sufficient number of positive full twists such that the resulting braid is positive, i.e. only positive crossings, see [12] for more details. The Euler-Floer characteristic remains unchanged. We denote a positive representative $x^+ \text{ rel } y^+$ again by $x \text{ rel } y$.

Define an augmented skeleton y^* by adding the constant strands $y_-(t) = (0, -1)$ and $y_+(t) = (0, 1)$. For proper braid classes it holds that $[x \text{ rel } y] = [x \text{ rel } y^*]$. For notational simplicity we denote the augmented skeleton again by y . We also choose the representative $x \text{ rel } y$ with the additional property that $\pi_2 x \text{ rel } \pi_2 y$ is a relative braid diagram, i.e. there are no tangencies between the strands, where π_2 the projection onto the q -coordinate. We denote the projection by $q \text{ rel } Q$, where $q = \pi_2 x$ and $Q = \pi_2 y$. Special braids on $\mathbb{R}^2$ can be constructed from (smooth) positive braids. Define $x_L = (q_t, q)$ and $y_L = (Q_t, Q)$, where the subscript t denotes differentiating with respect to t . These are called *Legendrian braids* with respect to $\theta = p dt - dq$.

Lemma 7.3. *For positive braid $x \text{ rel } y$ with only transverse, positive crossings, the braids $x_L \text{ rel } y_L$ and $x \text{ rel } y$ are isotopic as braids on $\mathbb{R}^2$. Moreover, if $x_L \text{ rel } y_L$ and $x'_L \text{ rel } y'_L$ are isotopic Legendrian braids, then they are isotopic via a Legendrian isotopy.*

Proof. By assumption $x \text{ rel } y$ is a representative for which the braid diagram $q \text{ rel } Q$ has only positive transverse crossings. Due to the transversality of intersections the associated Legendrian braid $x_L \text{ rel } y_L$ is a braid $[x \text{ rel } y]_{\mathbb{R}^2}$. Consider the homotopy

$$\zeta^j(t, \tau) = \tau p^j(t) + (1 - \tau) q_t^j,$$

for every strand q^j . At q -intersections, i.e. times t_0 such that $q^j(t_0) = q^{j'}(t_0)$ for some $j \neq j'$, it holds that $p^j(t_0) - p^{j'}(t_0)$ and $q_t^j(t_0) - q_t^{j'}(t_0)$ are non-zero and have the same sign since all crossings in $x \text{ rel } y$ are positive! Therefore, $\zeta^j(t_0, \tau) \neq \zeta^{j'}(t_0, \tau)$ for any intersection t_0 and any $\tau \in [0, 1]$, which shows that $x \text{ rel } y$ and $x_L \text{ rel } y_L$ are isotopic. Since $x_L \text{ rel } y_L$ and $x'_L \text{ rel } y'_L$ have only positive crossings, a smooth Legendrian isotopy exists. $\square$

The associated equivalence class of Legendrian braid diagrams is denoted by $[q \text{ rel } Q]$ and its fibers by $[q] \text{ rel } Q$.

7.3 Lagrangian systems

Legendrian braids can be described with Lagrangian systems and Hamiltonians of the form $H_L(x, t) = \frac{1}{2}p^2 - \frac{1}{2}q^2 + g(q, t)$. On the potential functions g we impose the following hypotheses:

(g1) $g \in C^\infty(\mathbb{R} \times \mathbb{R}/\mathbb{Z})$;

(g2) $\text{supp}(g) \subset [-R, R] \times \mathbb{R}/\mathbb{Z}$, for some $R > 1$.

In order to have a straightforward construction of a mechanical Lagrangian we may consider a special representation of y . The Euler-Floer characteristic $\chi(x \text{ rel } y)$ does not depend on the choice of the fiber $[x] \text{ rel } y$ and therefore also not on the skeleton y . We assume that y has linear crossings in y_L . Let $t = t_0$ be a crossing and let $I(t_0)$ be the set of labels defined by: $i, j \in I(t_0)$, if $i \neq j$ and $Q^i(t_0) = Q^j(t_0)$. A crossing at $t = t_0$ is *linear* if

$$Q_t^i = \text{constant}, \quad \forall i \in I(t_0), \quad \text{and} \quad \forall t \in (-\epsilon + t_0, \epsilon + t_0),$$

for some $\epsilon = \epsilon(t_0) > 0$. Every skeleton Q with transverse crossings is isotopic to a skeleton with linear crossings via a small local deformation at crossings. For Legendrian braids $x_L \text{ rel } y_L \in [x \text{ rel } y]_{\mathbb{R}^2}$ with linear crossings the following result holds:

Lemma 7.4. *Let y_L be a Legendrian skeleton with linear crossings. Then, there exists a Hamiltonian of the form $H_L(x, t) = \frac{1}{2}p^2 - \frac{1}{2}q^2 + g(q, t)$, with g satisfying Hypotheses (g1)-(g2), and $R > 0$ sufficiently large, such that y_L is a skeleton for $X_{H_L}(x, t)$.*

Proof. Due to the linear crossings in y_L we can follow the construction in [12]. For each strand Q^i we define the potentials $g^i(t, x) = -Q_{tt}^i(t)q$. By construction Q^i is a solution of the equation $Q_{tt}^i = -g_q^i(t, Q^i)$. Now choose small tubular neighborhoods of the strands Q^i and cut-off functions ω^i that are equal to 1 near Q^i and are supported in the tubular neighborhoods. If the tubular neighborhoods are narrow enough, then $\text{supp}(\omega^i g^i) \cap \text{supp}(\omega^j g^j) = \emptyset$, for all $i \neq j$, due to the fact that at crossings the functions g^i in question are zero. This implies that all strands Q^i satisfy the differential equation $Q_{tt}^i = -\sum_i \omega^i(t) g_q^i(Q^i, t)$ and on $[-1, 1] \times \mathbb{R}/\mathbb{Z}$, the function is $\sum_i \omega^i(t) g^i(q, t)$ is compactly supported. The latter follows from the fact that for the constant strands $Q^i = \pm 1$, the potentials g^i vanish. Let $R > 1$ and define

$$\tilde{g}^i(t, q) = \begin{cases} g^i(t, q) & \text{for } |q| \leq 1, t \in \mathbb{R}/\mathbb{Z}, \\ -\frac{1}{2m} q^2 & \text{for } |q| \geq R, t \in \mathbb{R}/\mathbb{Z}. \end{cases}$$

where $m = \#Q$, which yields smooth functions $\tilde{g}^i$ on $\mathbb{R} \times \mathbb{R}/\mathbb{Z}$. Now define

$$g(q, t) = \frac{1}{2} q^2 + \sum_{i=1}^m \tilde{g}^i(q, t).$$

By construction $\text{supp}(g) \subset [-R, R] \times \mathbb{R}/\mathbb{Z}$, for some $R > 1$ and the strands Q^i all satisfy the Euler-Lagrange equations $Q_{tt}^i = Q^i - g_{qq}(Q^i, t)$, which completes the proof. $\square$

The Hamiltonian H_L given by Lemma 7.4 gives rise to a Lagrangian system with the Lagrangian action given by

$$\mathcal{L}_g = \int_0^1 \frac{1}{2} \dot{q}_t^2 + \frac{1}{2} q^2 - g(q, t) dt. \quad (7.3)$$

The braid class $[q] \text{ rel } Q$ is bounded due to the special strands ± 1 and all free strands q satisfy $-1 \leq q(t) \leq 1$. Therefore, the set of critical points of $\mathcal{L}_g$ in $[q] \text{ rel } Q$ is a compact set. The critical points of $\mathcal{L}_g$ in $[q] \text{ rel } Q$ are in one-to-one correspondence with the zeroes of the equation

$$\Phi_{\mu, H_L}(x) = x - L_\mu^{-1}(\nabla H_L(x, t) + \mu x) = 0,$$

in the set $\Omega_{\mathbb{R}^2} = [x_L]_{\mathbb{R}^2} \text{ rel } y_L$, which implies that Φ_{μ, H_L} is a proper mapping on $\Omega_{\mathbb{R}^2}$. From Lemma 7.1 we derive that the zeroes of Φ_{μ, H_L} are contained in ball in $\mathbb{R}^2$ with radius $R' > 1$, and thus $\Phi_{\mu, H_L}^{-1}(0) \cap \Omega_{\mathbb{R}^2} \subset B_{R'}(0) \subset C^1(\mathbb{R}/\mathbb{Z})$. Therefore the Leray-Schauder degree is well-defined and in the generic case Lemma 4.2 and Equations (3.13), (4.6) and (5.2) yield

$$\deg_{LS}(\Phi_{\mu, H_L}, \Omega_{\mathbb{R}^2}, 0) = - \sum_{x \in \Phi_{\mu, H_L}^{-1}(0) \cap \Omega_{\mathbb{R}^2}} (-1)^{\mu^{CZ}(x)} = - \sum_{q \in \text{Crit}(\mathcal{L}_g) \cap ([q] \text{ rel } Q)} (-1)^{\beta(q)}. \quad (7.4)$$

We are now in a position to use a homotopy argument. We can scale y to a braid ρy such that the rescaled Legendrian braid ρy_L is supported in $\mathbb{D}^2$. By Lemma 7.3, y is isotopic to y_L and scaling defines an isotopy between y_L and ρy_L . Denote the isotopy from y to ρy_L by y_α . By Proposition 5.1 we obtain that for both skeletons y and ρy_L it holds that

$$\deg_{LS}(\Phi_{\mu,H}, \Omega, 0) = -\chi(x \text{ rel } y) = \deg_{LS}(\Phi_{\mu,H_\rho}, \Omega_\rho, 0),$$

where $\Omega_\rho = [\rho x_L] \text{ rel } \rho y_L \subset [x \text{ rel } y]$ and $H_\rho \in \mathcal{H}_\parallel(\rho y_L)$. Now extend H_ρ to $\mathbb{R}^2 \times \mathbb{R}/\mathbb{Z}$, such that Hypotheses (h1)-(h3) are satisfied for some $R > 1$. We denote the Hamiltonian again by H_ρ . By construction all zeroes of Φ_{μ,H_ρ} in $[\rho x_L] \text{ rel } \rho y_L$ are supported in $\mathbb{D}^2$ and therefore the zeroes of Φ_{μ,H_ρ} in $[\rho x_L]_{\mathbb{R}^2} \text{ rel } \rho y_L$ are also supported in $\mathbb{D}^2$. Indeed, any zero intersects $\mathbb{D}^2$, since the braid class is proper and since $\partial\mathbb{D}^2$ is invariant for the Hamiltonian vector field, a zero is either inside or outside $\mathbb{D}^2$. Combining these facts implies that a zero lies inside $\mathbb{D}^2$. This yields

$$\deg_{LS}(\Phi_{\mu,H_\rho}, \Omega_{\rho,\mathbb{R}^2}, 0) = \deg_{LS}(\Phi_{\mu,H_\rho}, \Omega_\rho, 0) = -\chi(x \text{ rel } y),$$

where $\Omega_{\rho,\mathbb{R}^2} = [\rho x_L]_{\mathbb{R}^2} \text{ rel } \rho y_L$. For the next homotopy we keep the skeleton ρy_L fixed as well as the domain $\Omega_{\rho,\mathbb{R}^2}$. Consider the linear homotopy of Hamiltonians

$$H_1(x, t; \alpha) = \frac{1}{2}p^2 - \frac{1}{2}q^2 + (1 - \alpha)h_\rho(x, t) + \alpha g_\rho(q, t),$$

where $H_{\rho,L}(t, x) = \frac{1}{2}p^2 - \frac{1}{2}q^2 + g_\rho(q, t)$ given by Lemma 7.4. This defines an admissible homotopy since ρy_L is a skeleton for all $\alpha \in [0, 1]$. The uniform estimates are obtained, as before, by Lemma 7.2, which allows application of the Leray-Schauder degree:

$$\deg_{LS}(\Phi_{\mu,H_{\rho,L}}, \Omega_{\rho,\mathbb{R}^2}, 0) = \deg_{LS}(\Phi_{\mu,H_\rho}, \Omega_{\rho,\mathbb{R}^2}, 0) = -\chi(x \text{ rel } y).$$

Finally, we scale ρy_L to y_L via $y_{\alpha,L} = (1 - \alpha)\rho y_L + \alpha y_L$ and we consider the homotopy

$$H_2(x, t; \alpha) = \frac{1}{2}p^2 - \frac{1}{2}q^2 + g(q, t; \alpha),$$

between H_L and $H_{\rho,L}$, where $g(q, t; \alpha)$ is found by applying Lemma 7.4 to $y_{\alpha,L}$. The uniform estimates from Lemma 7.2 allows us to apply the Leray-Schauder degree:

$$\deg_{LS}(\Phi_{\mu,H_L}, \Omega_{\mathbb{R}^2}, 0) = \deg_{LS}(\Phi_{\mu,H_{\rho,L}}, \Omega_{\rho,\mathbb{R}^2}, 0) = -\chi(x \text{ rel } y).$$

Combining the equalities for the various Leray-Schauder degrees with (7.4) yields:

$$-\deg_{LS}(\Phi_{H_L}, \Omega_{\mathbb{R}^2}, 0) = \chi(x \text{ rel } y) = \sum_{q \in \text{Crit}(\mathcal{L}_g) \cap ([q] \text{ rel } Q)} (-1)^{\beta(q)}. \quad (7.5)$$

7.4 Discretized braid classes

The Lagrangian problem (7.3) can be treated by using a variation on the method of broken geodesics. If we choose $1/d > 0$ sufficiently small, the integral

$$S_i(q_i, q_{i+1}) = \min_{\substack{q(t) \in E_i(q_i, q_{i+1}) \\ |q(t)| \leq 1}} \int_{\tau_i}^{\tau_{i+1}} \frac{1}{2} \dot{q}_t^2 + \frac{1}{2} q^2 - g(q, t) dt, \quad (7.6)$$

has a unique minimizer q^i , where $E_i(q_i, q_{i+1}) = \{q \in H^1(\tau_i, \tau_{i+1}) \mid q(\tau_i) = q_i, q(\tau_{i+1}) = q_{i+1}\}$, and $\tau_i = i/d$. Moreover, if $1/d$ is small, then the minimizers are non-degenerate and S_i is a smooth function of q_i and q_{i+1} . Critical points q of $\mathcal{L}_g$ with $|q(t)| \leq 1$ correspond to sequences $q_D = (q_0, \dots, q_d)$, with $q_0 = q_d$, which are critical points of the discrete action

$$\mathcal{W}(q_D) = \sum_{i=0}^{d-1} S_i(q_i, q_{i+1}). \quad (7.7)$$

A concatenation $\#_i q^i$ of minimizers q^i is continuous and is an element in the function space $H^1(\mathbb{R}/\mathbb{Z})$, and is referred to as a *broken geodesic*. The set of broken geodesics $\#_i q^i$ is denoted by $E(q_D)$ and standard arguments using the non-degeneracy of minimizers q^i show that $E(q_D) \hookrightarrow H^1(\mathbb{R}/\mathbb{Z})$ is a smooth, d -dimensional submanifold in $H^1(\mathbb{R}/\mathbb{Z})$. The submanifold $E(q_D)$ is parametrized by sequences $D_d = \{q_D \in \mathbb{R}^d \mid |q_i| \leq 1\}$ and yields the following commuting diagram:

$$\begin{array}{ccc} E(q_D) & \xrightarrow{\mathcal{L}_g} & \mathbb{R} \\ \#_i \downarrow & & \downarrow \mathcal{W} \\ & D_n & \end{array}$$

In the above diagram $\#_i$ is regarded as a mapping $q_D \mapsto \#_i q^i$, where the minimizers q_i are determined by q_D . The tangent space to $E(q_D)$ at a broken geodesic $\#_i q^i$ is identified by

$$\begin{aligned} T_{\#_i q^i} E(q_D) &= \{\psi \in H^1(\mathbb{R}/\mathbb{Z}) \mid -\psi_{tt} + \psi - g_{qq}(q^i(t), t)\psi = 0, \\ &\quad \psi(\tau_i) = \delta q_i, \quad \psi(\tau_{i+1}) = \delta q_{i+1}, \quad \delta q_i \in \mathbb{R}, \forall i\}, \end{aligned}$$

and $\#_i q^i + T_{\#_i q^i} E(q_D)$ is the tangent hyperplane at $\#_i q^i$. For $H^1(\mathbb{R}/\mathbb{Z})$ we have the following decomposition for any broken geodesic $\#_i q^i \in E(q_D)$:

$$H^1(\mathbb{R}/\mathbb{Z}) = E' \oplus T_{\#_i q^i} E(q_D), \quad (7.8)$$

where $E' = \{\eta \in H^1(\mathbb{R}/\mathbb{Z}) \mid \eta(\tau_i) = 0, \forall i\}$. To be more specific the decomposition is orthogonal with respect to the quadratic form

$$D^2 \mathcal{L}_g(q) \phi \tilde{\phi} = \int_0^1 \phi_t \tilde{\phi}_t + \phi \tilde{\phi} - g_{qq}(q(t), t) \phi \tilde{\phi} dt, \quad \phi, \tilde{\phi} \in H^1(\mathbb{R}/\mathbb{Z}).$$

Indeed, let $\eta \in E'$ and $\psi \in T_{\#_i q^i} E(q_D)$, then

$$\begin{aligned} D^2 \mathcal{L}_g(\#_i q^i) \eta \psi &= \sum_i \int_{\tau_i}^{\tau_{i+1}} \eta_t \psi_t + \eta \psi - g_{qq}(q^i(t), t) \xi \eta dt \\ &= \sum_i \psi_t \eta \Big|_{\tau_i}^{\tau_{i+1}} - \sum_i \int_{\tau_i}^{\tau_{i+1}} [-\psi_{tt} + \psi + g_{qq}(q^i(t), t) \psi] \eta dt = 0. \end{aligned}$$

Let $\phi = \eta + \psi$, then

$$D^2 \mathcal{L}_g(\#_i q^i) \phi \tilde{\phi} = D^2 \mathcal{L}_g(\#_i q^i) \eta \tilde{\eta} + D^2 \mathcal{L}_g(\#_i q^i) \psi \tilde{\psi},$$

by the above orthogonality. By construction the minimizers q^i are non-degenerate and therefore $D^2 \mathcal{L}_g|_{E'}$ is positive definite. This implies that the Morse index of a (stationary) broken geodesic is determined by $D^2 \mathcal{L}_g|_{T_{\#_i q^i} E(q_D)}$. By the commuting diagram for $\mathcal{W}$ this implies that the Morse index is given by quadratic form $D^2 \mathcal{W}(q_D)$. We have now proved the following lemma that relates the Morse index of critical points of the discrete action $\mathcal{W}$ to Morse index of the ‘full’ action $\mathcal{L}_g$.

Lemma 7.5. *Let q be a critical point of $\mathcal{L}_g$ and q_D the corresponding critical point of $\mathcal{W}$, then the Morse indices are the same i.e. $\beta(q) = \beta(q_D)$.*

For a 1-periodic function $q(t)$ we define the mapping

$$q \xrightarrow{D_d} q_D = (q_0, \dots, q_d), \quad q_i = q(i/d), \quad i = 0, \dots, d,$$

and q_D is called the discretization of q . The linear interpolation

$$q_D \mapsto \ell_{q_D}(t) = \#_i \left[q_i + \frac{q_{i+1} - q_i}{d} t \right],$$

reconstructs a piecewise linear 1-periodic function. For a relative braid diagram $q \text{ rel } Q$, let $q_D \text{ rel } Q_D$ be its discretization, where Q_D is obtained by applying D_d to every strand in Q . A discretization $q_D \text{ rel } Q_D$ is *admissible* if $\ell_{q_D} \text{ rel } \ell_{Q_D}$ is homotopic to $q \text{ rel } Q$, i.e. $\ell_{q_D} \text{ rel } \ell_{Q_D} \in [q \text{ rel } Q]$. Define the *discrete* relative braid class $[q_D \text{ rel } Q_D]$ as the set of ‘discrete relative braids’ $q'_D \text{ rel } Q'_D$, such that $\ell_{q'_D} \text{ rel } \ell_{Q'_D} \in [q \text{ rel } Q]$. The associated fibers are denoted by $[q_D] \text{ rel } Q_D$. It follows from [8], Proposition 27, that $[q_D \text{ rel } Q_D]$ is guaranteed to be connected when

$$d > \#\{\text{crossings in } q \text{ rel } Q\},$$

i.e. for any two discrete relative braids $q_D \text{ rel } Q_D$ and $q'_D \text{ rel } Q'_D$, there exists a homotopy $q_D^\alpha \text{ rel } Q_D^\alpha$ (discrete homotopy) such that $\ell_{q_D^\alpha} \text{ rel } \ell_{Q_D^\alpha}$ is a path in $[q \text{ rel } Q]$. Note that fibers are not necessarily connected! For a braid classes $[q \text{ rel } Q]$ the associated discrete braid class $[q_D \text{ rel } Q_D]$ may be connected for a smaller choice of d .

We showed above that if $1/d > 0$ is sufficiently small, then the critical points of $\mathcal{L}_g$, with $|q| \leq 1$, are in one-to-one correspondence with the critical points of $\mathcal{W}$, and their Morse indices coincide by Lemma 7.5. Moreover, if $1/d > 0$ is small enough, then for all critical points of $\mathcal{L}_g$ in $[q] \text{ rel } Q$, the associated discretizations are admissible and $[q_D] \text{ rel } Q_D$ is a connected set. The discretizations of the critical points of $\mathcal{L}_g$ in $[q] \text{ rel } Q$ are critical points of $\mathcal{W}$ in the discrete braid class fiber $[q_D] \text{ rel } Q_D$.

Now combine the index identity with (7.5), which yields

$$\chi(x \text{ rel } y) = \sum_{q \in \text{Crit}(\mathcal{L}_g) \cap ([q] \text{ rel } Q)} (-1)^{\beta(q)} = \sum_{q_D \in \text{Crit}(\mathcal{W}) \cap ([q_D] \text{ rel } Q_D)} (-1)^{\beta(q_D)}. \quad (7.9)$$

7.5 The Conley index for discrete braids

In [8] an invariant for discrete braid classes $[q_D] \text{ rel } Q_D$ is defined based on the Conley index. The invariant $\text{HC}_*([q_D] \text{ rel } Q_D)$ is independent of the fiber and can be described as follows. A fiber $[q_D] \text{ rel } Q_D$ is a finite dimensional cube complex with a finite number of connected components. Denote the closures of the connected components by N_j . The faces of the hypercubes N_j can be co-oriented in direction of decreasing the number of crossing in $q_D \text{ rel } Q_D$, and define N_j^- as the closure of the set of faces with outward pointing co-orientation. The sets N_j^- are called *exit sets*. The invariant is given by

$$\text{HC}_*([q_D] \text{ rel } Q_D) = \bigoplus_j H_*(N_j, N_j^-).$$

The invariant is well-defined for any $d > 0$ for which there exist admissible discretizations and is independent of both the fiber and the discretization size. From [8] we have for any Morse function $\mathcal{W}$ on a proper braid class fiber $[q_D] \text{ rel } Q_D$,

$$\sum_{q_D \in \text{Crit}(\mathcal{W}) \cap ([q_D] \text{ rel } Q_D)} (-1)^{\beta(q_D)} = \chi(\text{HC}_*([q_D] \text{ rel } Q_D)) =: \chi(q_D \text{ rel } Q_D). \quad (7.10)$$

The latter can be computed for any admissible discretization and is an invariant for $[q \text{ rel } Q]$. Combining 7.9 and 7.10 gives

$$\chi(x \text{ rel } y) = \chi(q_D \text{ rel } Q_D). \quad (7.11)$$

In this section we assumed without loss of generality that $x \text{ rel } y$ is augmented and since the Euler-Floer characteristic is a braid class invariant, an admissible discretization is construction for an appropriate augmented, Legendrian representative $x_L \text{ rel } y_L$. Summarizing

$$\chi(x \text{ rel } y) = \chi(x_L \text{ rel } y_L^*) = \chi(q_D \text{ rel } Q_D^*).$$

Since $\chi(q_D \text{ rel } Q_D^*)$ is the same for any admissible discretization, the Euler-Floer characteristic can be computed using any admissible discretization, which proves Theorem 1.3.

Remark 7.1. The invariant $\chi(q_D \text{ rel } Q_D)$ is a true Euler characteristic of a topological pair. To be more precise

$$\chi(q_D \text{ rel } Q_D) = \chi([q_D] \text{ rel } Q_D, [q_D]^- \text{ rel } Q_D),$$

where $[q_D]^- \text{ rel } Q_D$ is the exit set as described above. A similar characterization does not a priori exist for $[x] \text{ rel } y$. Firstly, it is more complicated to designate the equivalent of an exit set $[x]^- \text{ rel } y$ for $[x] \text{ rel } y$, and secondly it is not straightforward to develop a (co)-homology theory that is able to provide meaningful information about the topological pair $([x] \text{ rel } y, [x]^- \text{ rel } y)$. This problem is circumvented by considering Hamiltonian systems and carrying out Floer's approach towards Morse theory (see [7]), by using the isolation property of $[x] \text{ rel } y$. The fact that the Euler characteristic of Floer homology is related to the Euler characteristic of topological pair indicates that Floer homology is a good substitute for a suitable (co)-homology theory.

8 Examples

We will illustrate by means of two examples that the Euler-Floer characteristic is computable and can be used to find closed integral curves of vector fields on the 2-disc.

8.1 Example

Figure 1[left] shows the braid diagram $q \text{ rel } Q$ of a positive relative braid $x \text{ rel } y$. The discretization with $q_D \text{ rel } Q_D$, with $d = 2$, is shown in Figure 1[right]. The chosen discretization is admissible and defines the relative braid class $[q_D \text{ rel } Q_D]$. There are five strands, one is free and four are fixed. We denote the points on the free strand by $q_D = (q_0, q_1)$ and on the skeleton by $Q_D = \{Q^1, \dots, Q^4\}$, with $Q^i = (Q_0^i, Q_1^i)$, $i = 1, \dots, 4$.

In Figure 2[left] the braid class fiber $[q_D] \text{ rel } Q_D$ is depicted. The coordinate q_0 is allowed to move between Q_0^3 and Q_0^2 and q_1 remains in the same braid class if it varies between Q_1^1 and Q_1^4 . For the values $q_0 = Q_0^3$ and $q_0 = Q_0^2$ the relative braid becomes singular and if q_0 crosses these values two intersections are created. If q_1




Figure 1: A positive braid diagram [left] and an admissible discretization [right].

crosses the values Q_1^1 or Q_1^4 two intersections are destroyed. This provides the desired co-orientation, see Figure 2[middle]. The braid class fiber $[q_D] \text{ rel } Q_D$ consists of 1 component and we have that

$$N = \text{cl}([q_D] \text{ rel } Q_D) = \{(q_0, q_1) : Q_0^3 \leq q_0 \leq Q_0^2, Q_1^1 \leq q_1 \leq Q_1^4\},$$

and the exit set is

$$N^- = \{(q_0, q_1) : q_1 = Q_1^1, \text{ or } q_1 = Q_1^4\}.$$

For the Conley index this gives:

$$\text{HC}_k([q_D] \text{ rel } Q_D) = H_k(N, N^-; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & k = 1 \\ 0 & \text{otherwise} \end{cases}$$




Figure 2: The relative braid fiber $[q_D] \text{ rel } Q_D$ and $N = \text{cl}([q_D] \text{ rel } Q_D)$.

The Euler characteristic of $([q_D] \text{ rel } Q_D, [q_D]^- \text{ rel } Q_D)$ can be computed now and the Euler-Floer characteristic $(x \text{ rel } y)$ is given by

$$\chi(x \text{ rel } y) = \chi([q_D] \text{ rel } Q_D, [q_D]^- \text{ rel } Q_D) = -1 \neq 0$$

From Theorem 1.2 we derive that any vector field for which y is a skeleton has at least 1 closed integral curve $x_0 \text{ rel } y \in [x] \text{ rel } y$. Theorem 1.2 also implies that any orientation preserving diffeomorphism f on the 2-disc which fixes the set of four points A_4 , whose mapping class $[f; A_4]$ is represented by the braid y has an additional fixed point.

8.2 Example

The theory can also be used to find additional closed integral curves by concatenating the skeleton y . As in the previous example y is given by Figure 1. Glue ℓ copies of the skeleton y to its ℓ -fold concatenation and a reparametrize time by $t \mapsto \ell \cdot t$. Denote the rescaled ℓ -fold concatenation of y by $\#_\ell y$. Choose $d = 2\ell$ and discretize $\#_\ell y$ as in the previous example. For a given braid class $[x \text{ rel } \#_\ell y]$, Figure 3 below




Figure 3: A discretization of a braid class with a 5-fold concatenation of the skeleton y . The number of odd anchor points in middle position is $\mu = 3$.

shows a discretized representative $q_D \text{ rel } \#_\ell Q_D$, which is admissible. For the skeleton $\#_\ell Q_D$ we can construct $3^\ell - 2$ proper relative braid classes in the following way: the even anchor points of the free strand q_D are always in the middle and for the odd anchor points we have 3 possible choices: bottom, middle, top (2 braids are not proper). We now compute the Conley index of the $3^\ell - 2$ different proper discrete relative braid classes and show that the Euler-Floer characteristic is non-trivial for these relative braid classes.

The configuration space $N = \text{cl}([q_D] \text{ rel } \#_\ell Q_D)$ in this case is given by a cartesian product of 2ℓ closed intervals, and therefore a 2ℓ -dimensional hypercube. We now proceed by determining the exit set N^- . As in the previous example the co-orientation is found by a union of faces with an outward pointing co-orientation.

Due to the simple product structure of N , the set N^- is determined by the odd anchor points in the middle position. Denote the number of middle positions at odd anchor points by μ . In this way N^- consists of opposite faces at odd anchor points in middle position, see Figure 3. Therefore

$$\mathrm{HC}_k([q_D] \operatorname{rel} \#_\ell Q_D) = H_k(N, N^-) = \begin{cases} \mathbb{Z}_2 & k = \mu \\ 0 & k \neq \mu, \end{cases}$$

and the Euler-Floer characterisc is given by

$$\chi(x \operatorname{rel} \#_\ell y) = (-1)^\mu \neq 0.$$

Let $X(x, t)$ be a vector field for which y is a skeleton of closed integral curves, then $\#_\ell y$ is a skeleton for the vector field $X^\ell(x, t) := \ell X_\ell(x, \ell t)$. From Theorem 1.2 we derive that there exists a closed integral curve in each of the $3^\ell - 2$ proper relative classes $[x] \operatorname{rel} y$ described above. For the original vector field X this yields $3^\ell - 2$ distinct closed integral curves. Using the arguments in [13] one can find a compact invariant set for X with positive topological entropy, which proves that the associated flow is ‘chaotic’ whenever y is a skeleton of given integral curves

8.3 Example

So far we have not addressed the question whether the closed integral curves $x \operatorname{rel} y$ are non-trivial, i.e. not equilibrium points of X . The theory can also be extended in order to find non-trivial closed integral curves. This paper restricts to relative braids where x consists of just one strand. Braid Floer homology for relative braids with x consisting of n strands is defined in [12]. To illustrate the importance of multi-strand braids we consider the discrete braid class in Figure 4.

The braid class depicted in Figure 4[right] is discussed in the previous example and the Euler-Floer characteristic is equal to 1. By considering all translates of x on the circle $\mathbb{R}/\mathbb{Z}$, we obtain the braids in Figure 4[left]. The latter braid class is proper and encodes extra information about q_D relative to Q_D . The braid class fiber is a 6-dimensional cube with the same Conley index as the braid class in Figure 4[right]. Therefore,

$$\chi(q_D \operatorname{rel} Q_D) = (-1)^2 = 1.$$

As in the 1-strand case, the discrete Euler characteristic can be used to compute the associated Euler-Floer characteristic of $x \operatorname{rel} y$ and $\chi(x \operatorname{rel} y) = 1$. The skeleton y thus forces solutions $x \operatorname{rel} y$ of the above described type. The additional information we obtain this way is that for braid classes $[x \operatorname{rel} y]$, the associated closed integral curves for X cannot be constant and therefore represent non-trivial closed integral curves.




Figure 4: A discretization of a braid class with a 3-fold concatenation of the skeleton y . The number of odd anchor points in middle position is $\mu = 2$ [right]. If we represent all translates of x we obtain a proper relative braid class where x is a 3-strand braid [left]. The latter provides additional linking information.

References

- [1] A Abbondandolo, *On the Morse index of Lagrangian systems*, Nonlinear Analysis **53** (2003), no. 3-4, 551–566.
- [2] J. J. Duistermaat, *On the Morse index in variational calculus*, Advances in Mathematics **21** (1976), no. 2, 173–195.
- [3] P.M. Fitzpatrick and J. Pejsachowicz, *The fundamental group of the space of linear Fredholm operators and the global analysis of semilinear equations*, Fixed point theory and its applications (Berkeley, CA, 1986), Contemp. Math., vol. 72, Amer. Math. Soc., Providence, RI, 1988, pp. 47–87. MR 956479 (89h:47097)
- [4] ———, *Parity and generalized multiplicity*, Trans. Amer. Math. Soc. **326** (1991), no. 1, 281–305. MR 1030507 (91j:58038)
- [5] P.M. Fitzpatrick, J. Pejsachowicz, and P. Rabier, *The degree of proper c^2 -fredholm mappings*, Journal für die Reine und Angewandte Mathematik **427** (1992), 1–33.
- [6] P.M. Fitzpatrick, J. Pejsachowicz, and L. Recht, *Spectral flow and bifurcation of critical points of strongly-indefinite functionals. i. general theory*, Journal of Functional Analysis **162** (1999), no. 1, 52–95.
- [7] A. Floer, *Symplectic fixed points and holomorphic spheres*, Comm. Math. Phys. **120** (1989), no. 4, 575–611. MR MR987770 (90e:58047)
- [8] R. Ghrist, J.B. van den Berg, and R.C. Vandervorst, *Morse theory on spaces of braids and lagrangian dynamics*, Invent. Math. **152** (2003), no. 2, 369–432.

- [9] N. G. Lloyd, *Degree theory*, Cambridge University Press, Cambridge, 1978, Cambridge Tracts in Mathematics, No. 73. MR MR0493564 (58 #12558)
- [10] J. Robbin and D. Salamon, *The Maslov index for paths*, *Topology* **32** (1993), no. 4, 827–844. MR MR1241874 (94i:58071)
- [11] ———, *The spectral flow and the Maslov index*, *Bull. London Math. Soc.* **27** (1995), 1–33.
- [12] J.B. van den Berg, R. Ghrist, R.C. Vandervorst, and W. Wójcik, *Braid Floer homology*, preprint arXiv:0910.0647 (2010), 1–55.
- [13] J.B. van den Berg, R.C. Vandervorst, and W. Wójcik, *Chaos in orientation reversing twist maps on the plane*, *Topology Appl.* **154** (2007), 2580–2606.